

Left Quotients of a C*-algebra, I: Representation via vector sections *

Ngai-Ching Wong

Department of Applied Mathematics

National Sun Yat-sen University

Kao-hsiung, 80424, Taiwan, R.O.C.

E-mail: wong@math.nsysu.edu.tw

Abstract

Let A be a C*-algebra, L a closed left ideal of A and p the closed projection related to L . We show that for an xp in $A^{**}p$ ($\cong A^{**}/L^{**}$) if $pAxp \subset pAp$ and $px^*xp \in pAp$ then $xp \in Ap$ ($\cong A/L$). The proof goes by interpreting elements of $A^{**}p$ (resp. Ap) as admissible (resp. continuous admissible) vector sections over the base space $F(p) = \{\varphi \in A^* : \varphi \geq 0, \varphi(p) = \|\varphi\| \leq 1\}$ in the notions developed by Dixmier and Douady, Fell, and Tomita. We consider that our results complement both Kadison function representation and Takesaki duality theorem.

1 Introduction

It is known that every closed left ideal L of a C*-algebra A is related to a closed projection p in the sense that $L = A^{**}(1-p) \cap A$ (and thus $L^{**} = A^{**}(1-p)$). Moreover, A/L (resp. A^{**}/L^{**}) is isometrically isomorphic to Ap (resp. $A^{**}p$) as Banach spaces [?, ?, ?]. Here, a projection p in A^{**} is said to be *closed* if the face $F(p) = \{\varphi \in Q(A) : \varphi(p) = \|\varphi\|\}$ of the weak* compact convex set $Q(A) = \{\varphi \in A^* : \varphi \geq 0, \|\varphi\| \leq 1\}$ is closed (cf. [?]).

*This research is partially supported by National Science Council of Taiwan, R.O.C., NSC 81-0208-M-110-512.

Keywords: C*-algebras, continuous fields of Hilbert spaces, left quotients, affine property.

1991 Mathematics Subject Classification. 46L05, 46L45.

For each φ in $F(p)$, $L_\varphi = \{a \in A : \varphi(a^*a) = 0\}$ is a closed left ideal of A , and $L = \bigcap_{\varphi \in F(p)} L_\varphi$. This gives a natural embedding

$$A/L \hookrightarrow \prod_{\varphi \in F(p)} A/L_\varphi, \quad a + L \longmapsto (a + L_\varphi)_{\varphi \in F(p)}.$$

Let H_φ be the completion of the pre-Hilbert space A/L_φ with respect to the inner product $\langle a + L_\varphi, b + L_\varphi \rangle_\varphi := \varphi(b^*a)$ for each φ in $F(p)$ (*i.e.* the GNS construction for φ). In this way, $A/L \cong Ap$ is embedded into the field of Hilbert spaces $(F(p), \{H_\varphi\}_\varphi)$. This also induces an embedding of $A^{**}/L^{**} \cong A^{**}p$ into $(F(p), \{H_\varphi\}_\varphi)$, since we can identify H_φ with the GNS Hilbert space for φ when φ is regarded as a positive functional on A^{**} and the GNS representation of A^{**} extends that of A .

By a result of Brown [?, 3.5], pAp (resp. $pA^{**}p$) is isometrically order isomorphic to the Banach space of all continuous (resp. bounded) affine functions on $F(p)$ which vanish at zero. In particular, for every xp in Ap the scalar maps $\varphi \longmapsto \varphi(pa^*xp) = \varphi(a^*x)$, $\forall a \in A$, and $\varphi \longmapsto \varphi(px^*xp) = \varphi(x^*x)$ are continuous on $F(p)$. In this paper, we proved that if xp in $A^{**}p$ satisfies conditions that the scalar maps $\varphi \longmapsto \varphi(a^*x)$, $\forall a \in A$, and $\varphi \longmapsto \varphi(x^*x)$ are continuous on $F(p)$ then $xp \in Ap$. In other words, for xp in $A^{**}p$, $pAxp \subset pAp$ and $px^*xp \in pAp$ imply $xp \in Ap$.

We establish the above result by first looking for an admissibility condition characterizing those vector sections of the field of Hilbert spaces $(F(p), \{H_\varphi\}_\varphi)$ arising from elements of $A^{**}p$ (theorem ??). Then, following ideas of Fell [?] and Dixmier and Douady [?], we implement a continuous structure $\Gamma(Ap)$ of $(F(p), \{H_\varphi\}_\varphi)$ in which all vector sections arising from elements of Ap are continuous. Finally, we prove that continuous admissible vector sections of $(F(p), \{H_\varphi\}_\varphi, \Gamma(Ap))$ are exactly those arising from elements of Ap (theorem ??). And this is translated to the result just mentioned above (corollary ??).

The way we look at elements of $A^{**}p$ and Ap as admissible vector sections and continuous admissible vector sections over the compact convex set $F(p)$ suggests some interesting questions and results. For example, it is natural to ask for an xp in $A^{**}p$ if the continuity of the scalar maps $\varphi \longmapsto \varphi(a^*x)$, $\forall a \in A$, and $\varphi \longmapsto \varphi(x^*x)$ on the extreme boundary $F(p) \cap (P(A) \cup \{0\})$ of $F(p)$ can imply $xp \in Ap$, where $P(A)$ is the pure state space of A . In [?], we prove that such an xp has a continuous atomic part in many cases, *i.e.* there is an ap in Ap such that $zxp = zap$, where z is the maximal atomic projection of A . Even when $p = 1$, this is new and supplements results of Shultz [?] and Brown [?], which say that for an x in A^{**} if $\varphi \longmapsto \varphi(x)$ is uniformly continuous on $P(A) \cup \{0\}$ then $zx \in zA$. On the other hand, following the plan of Tomita [?] and

using ideas of Rieffel [?], we represent bounded Banach space operators on $A/L \cong Ap$ as fields of bounded Hilbert space operators in the context of $(F(p), \{H_\varphi\}_\varphi, \Gamma(Ap))$. Many ideas of Tomita about the theory of left regular representation of A on A/L can thus be implemented in this context (see [?]).

When $p = 1$, one can easily find the origin of our theory from Kadison function representation (see section ??) and Takesaki duality theorem [?, ?, ?] (see section ??). However, the results of Kadison and Takesaki are not ready to apply to left quotients $Ap (\cong A/L)$ if $p \neq 1$ (*i.e.* $L \neq \{0\}$). To extend these classical tools to the general case of $p \neq 1$ as shown in this paper, Tomita [?, ?] indicates us a way to set up our theory and Akemann [?, ?, ?], Dixmier and Douady [?], Effros [?], Fell [?], and Prosser [?] provide us the basic machinery.

We would like to express our deep gratitude to Professor L.G. Brown for many valuable advices. This paper is based on the author's doctoral dissertation [?] under his supervision.

2 Represent W^* -algebras via admissible vector sections

Let M be a W^* -algebra with predual M_* and $Q_*(M) = \{\varphi \in M_* : \varphi \geq 0, \|\varphi\| \leq 1\}$. Let p be a projection in M and $F(p) = \{\varphi \in Q_*(M) : \varphi(p) = \|\varphi\|\}$, the face of the convex set $Q_*(M)$ supported by p . For each φ in $F(p)$, the GNS construction yields a cyclic representation $(\pi_\varphi, H_\varphi, \omega_\varphi)$ of M . $\overline{\pi_\varphi(M)\omega_\varphi} = H_\varphi$ and $\varphi(x) = \langle \pi_\varphi(x)\omega_\varphi, \omega_\varphi \rangle_\varphi, \forall x \in M$, where $\langle \cdot, \cdot \rangle_\varphi$ is the inner product of the Hilbert space H_φ . Write $x\omega_\varphi$ for $\pi_\varphi(x)\omega_\varphi, \forall x \in M, \forall \varphi \in F(p)$. Note that $p\omega_\varphi = \omega_\varphi, \forall \varphi \in F(p)$. By convention, we set H_φ to be the zero dimensional Hilbert space when $\varphi = 0$. In this way, there is an embedding $Mp \hookrightarrow \prod_{\varphi \in F(p)} H_\varphi$ defined by $xp \longmapsto (x\omega_\varphi)_{\varphi \in F(p)}$. If we equip the range of this embedding with the ℓ^∞ norm then it is even an isometry as

$$\|xp\|^2 = \sup_{\varphi \in Q_*(M)} \varphi(pxp) = \sup_{\varphi \in F(p)} \varphi(x^*x) = \sup_{\varphi \in F(p)} \|x\omega_\varphi\|_\varphi^2 = \|(x\omega_\varphi)_{\varphi \in F(p)}\|_\infty^2.$$

We are going to classify those vector sections in $\prod_{\varphi \in F(p)} H_\varphi$ arising from this embedding. First, we observe that fibers H_φ in $\prod_{\varphi \in F(p)} H_\varphi$ are not independent of each other. The following definition is taken from Tomita [?] (with a slight modification).

Definition 2.1 Let M be a W^* -algebra. For each ψ in M_* and each φ in $F(p)$, we set

$$\|\psi\|_\varphi = \sup\{|\psi(x)| : x \in M \text{ and } \|x\omega_\varphi\|_\varphi = \varphi(x^*x)^{1/2} \leq 1\},$$

and $L^2(\varphi) = \{\psi \in M_* : \|\psi\|_\varphi < \infty\}$. We say that ψ is *observable at φ* if $\psi \in L^2(\varphi)$. It follows from the Riesz–Fréchet theorem that for each ψ in $L^2(\varphi)$ there is a unique $\omega_{\psi\varphi}$ in H_φ such that

$$\psi(x) = \langle x\omega_\varphi, \omega_{\psi\varphi} \rangle_\varphi, \quad \forall x \in M.$$

It can be verified that the map Λ defined by $\Lambda_\varphi(\psi) = \omega_{\psi\varphi}$ is a conjugate isometrical isomorphism from $L^2(\varphi)$ onto H_φ [?]. The proof of the following lemma is left to the readers.

Lemma 2.2 *For each ψ in $L^2(\varphi)$ and x in M , we have*

$$\Lambda_\varphi(x\psi) = x^* \Lambda_\varphi(\psi).$$

In other words,

$$\omega_{(x\psi)\varphi} = x^* \omega_{\psi\varphi},$$

where $x\psi$ in $MF(p)$ is defined by $x\psi(y) = \psi(xy), \forall y \in M$.

Definition 2.3 For each ψ, φ in $F(p)$ with $0 \leq \psi \leq \lambda\varphi$ for some $\lambda > 0$, let

$$T_{\psi\varphi} : H_\varphi \longrightarrow H_\psi$$

be the linear map from H_φ into H_ψ sending $x\omega_\varphi$ to $x\omega_\psi$. Note that $\psi \in L^2(\varphi)$ by the Cauchy–Schwartz inequality. Moreover, $\|T_{\psi\varphi}\| \leq \lambda^{1/2}$ and $T_{\psi\varphi}^*(x\omega_\psi) = x\omega_{\psi\varphi} = \Lambda_\varphi(x^*\psi), \forall x \in M$.

Definition 2.4 A vector section $f : \varphi \longmapsto f(\varphi) \in H_\varphi$ is said to be *admissible* over $F(p)$ if whenever $\psi, \varphi \in F(p)$ such that $0 \leq \psi \leq \varphi$,

$$T_{\psi\varphi}(f(\varphi)) = f(\psi).$$

f is said to be an *affine vector section* over $F(p)$ if the functional

$$\varphi \longmapsto \langle f(\varphi), x\omega_\varphi \rangle_\varphi$$

is affine on the convex set $F(p)$ for each x in M .

It is easy to see that whenever $0 \leq \psi \leq \varphi \leq \rho$ in $F(p)$, $T_{\psi\varphi}T_{\varphi\rho} = T_{\psi\rho}$. Moreover, for an admissible vector section f and φ, ψ in $F(p)$ such that $0 \leq \psi \leq \lambda\varphi$ for some $\lambda > 0$, we have $T_{\psi\varphi}f(\varphi) = f(\psi)$, too.

Proposition 2.5 *Every admissible vector section $f = (f(\varphi))_\varphi$ over $F(p)$ is bounded, i.e. $\|f\|_\infty = \sup_{\varphi \in F(p)} \|f(\varphi)\|_\varphi < \infty$.*

PROOF. Assume the contrary and choose φ_n in $F(p)$ such that

$$\|f(\varphi_n)\|_{\varphi_n} > 2^n, n = 1, 2, \dots$$

Set

$$\varphi = \sum_n \frac{1}{2^n} \varphi_n$$

in $F(p)$. Since $0 \leq \varphi_n \leq 2^n \varphi$, $T_{\varphi_n \varphi}$ in $B(H_\varphi, H_{\varphi_n})$ exists and $\|T_{\varphi_n \varphi}\|^2 \leq 2^n$. Therefore,

$$\|f(\varphi_n)\|_{\varphi_n}^2 = \|T_{\varphi_n \varphi} f(\varphi)\|_{\varphi_n}^2 \leq 2^n \|f(\varphi)\|_\varphi^2, n = 1, 2, \dots$$

Hence

$$\|f(\varphi)\|_\varphi^2 \geq \frac{1}{2^n} \|f(\varphi_n)\|_{\varphi_n}^2 \geq \frac{1}{2^n} 2^{2n} = 2^n, n = 1, 2, \dots,$$

a contradiction. □

Here is our main result:

Theorem 2.6 *Mp is isometrically isomorphic to the Banach space of all admissible vector sections in $\prod_{\varphi \in F(p)} H_\varphi$ equipped with the norm $\|\cdot\|_\infty$. A vector section in $\prod_{\varphi \in F(p)} H_\varphi$ is admissible if and only if it is bounded and affine.*

We shall present the proof of theorem ?? in two parts. It is trivial that each element of Mp defines an admissible vector section over $F(p)$ in the manner described at the start of this section. For the converse, we shall associate to each admissible vector section $f = (f(\varphi))_{\varphi \in F(p)}$ a bounded linear functional \tilde{f} of the predual $\{\varphi(\cdot p)|_{Mp} : \varphi \in M_*\}$ of Mp , which can be identified with $MF(p) = \{x\varphi(\cdot) = \varphi(x\cdot) : x \in M, \varphi \in F(p)\}$ (cf. [?]).

Lemma 2.7 *Let $f = (f(\varphi))_\varphi$ be an admissible vector section over $F(p)$.*

(a) *If ϕ in M_* is observable at both φ and ψ in $F(p)$ then*

$$\langle f(\varphi), \omega_{\phi\varphi} \rangle_\varphi = \langle f(\psi), \omega_{\phi\psi} \rangle_\psi.$$

(b) If $\phi \in M_*$ and $\varphi, \psi \in F(p)$ such that

$$\phi = x^* \varphi = y^* \psi$$

for some x, y in M then

$$\langle f(\varphi), \omega_{\phi\varphi} \rangle_{\varphi} = \langle f(\varphi), x\omega_{\varphi} \rangle_{\varphi} = \langle f(\psi), y\omega_{\psi} \rangle_{\psi} = \langle f(\psi), \omega_{\phi\psi} \rangle_{\psi}.$$

PROOF. (a) Let $\rho = \frac{\varphi + \psi}{2} \in F(p)$. By the admissibility of f , $T_{\varphi\rho}(f(\rho)) = f(\varphi)$ and $T_{\psi\rho}(f(\rho)) = f(\psi)$. It is easy to see that $T_{\varphi\rho}^*(\omega_{\phi\varphi}) = \omega_{\phi\rho}$ and $T_{\psi\rho}^*(\omega_{\phi\psi}) = \omega_{\phi\rho}$. Now

$$\langle f(\varphi), \omega_{\phi\varphi} \rangle_{\varphi} = \langle T_{\varphi\rho} f(\rho), \omega_{\phi\varphi} \rangle_{\varphi} = \langle f(\rho), T_{\varphi\rho}^*(\omega_{\phi\varphi}) \rangle_{\rho} = \langle f(\rho), \omega_{\phi\rho} \rangle_{\rho} = \langle f(\psi), \omega_{\phi\psi} \rangle_{\psi}.$$

(b) First, note that ϕ is observable at both φ and ψ and thus the asserted equalities make sense. Our assertion follows from lemma ?? and (a) and the following observation

$$\omega_{\phi\varphi} = \Lambda_{\varphi}(\phi) = \Lambda_{\varphi}(x^* \varphi) = x\Lambda_{\varphi}(\varphi) = x\omega_{\varphi}$$

and

$$\omega_{\phi\psi} = \Lambda_{\psi}(\phi) = \Lambda_{\psi}(y^* \psi) = y\Lambda_{\psi}(\psi) = y\omega_{\psi}.$$

□

Note that $Mp = (MF(p))^*$. This suggests us to make the following

Definition 2.8 Let $f = (f(\varphi))_{\varphi}$ be an admissible vector section over $F(p)$. Define for each ϕ in $MF(p)$,

$$\tilde{f}(\phi) = \langle f(\varphi), \omega_{\phi\varphi} \rangle_{\varphi},$$

where $\varphi \in F(p)$ and ϕ is observable at φ .

Clearly, $\tilde{f}(0) = 0$. For a non-zero ϕ in $MF(p)$, it follows from lemma ??(a) that the definition of $\tilde{f}(\phi)$ is independent of the choice of φ for which $\phi \in L^2(\varphi)$, and $\varphi = |\phi|/\|\phi\|$ is just a good choice, where $|\phi|$ is the absolute value of ϕ coming from the polar decomposition of ϕ (see, e.g. [?]). Moreover, if $\phi = x^* \varphi$ for some φ in $F(p)$ then by lemma ??(b),

$$\tilde{f}(\phi) = \langle f(\varphi), x\omega_{\varphi} \rangle_{\varphi}.$$

PROOF OF THE FIRST PART OF THEOREM ??. The first task is to prove that \tilde{f} is a bounded linear functional of $MF(p)$ for every admissible vector section $f = (f(\varphi))_{\varphi}$ over $F(p)$. To verify that \tilde{f} is additive, let ρ, φ and ψ be elements of $MF(p)$ such that

$\rho = \varphi + \psi$. In case $\varphi = \psi = 0$, it is plain that $\tilde{f}(\rho) = \tilde{f}(\varphi) + \tilde{f}(\psi)$. Suppose that not both φ and ψ are zero. [?] or [?] showed that

$$\|\rho(x)\|^2 \leq (\|\varphi\| + \|\psi\|)(|\varphi| + |\psi|)(x^*x), \quad \forall x \in M.$$

Hence, $|\rho| \in L^2(\tau)$, where $\tau = \frac{|\varphi|+|\psi|}{\|\varphi\|+\|\psi\|} \in F(p)$. Clearly, $|\varphi|, |\psi| \in L^2(\tau)$. As a result, ρ, φ and $\psi \in L^2(\tau)$. Now, $\tilde{f}(\rho) = \langle f(\tau), \omega_{\rho\tau} \rangle_\tau$, $\tilde{f}(\varphi) = \langle f(\tau), \omega_{\varphi\tau} \rangle_\tau$ and $\tilde{f}(\psi) = \langle f(\tau), \omega_{\psi\tau} \rangle_\tau$. The additivity of \tilde{f} follows easily since, by uniqueness, $\omega_{\rho\tau} = \omega_{\varphi\tau} + \omega_{\psi\tau}$. By lemma ??, $\omega_{(\lambda\phi)\varphi} = \bar{\lambda}\omega_{\phi\varphi}$, $\forall \lambda \in \mathbb{C}, \forall \varphi \in F(p), \forall \phi \in L^2(\varphi)$. Therefore, \tilde{f} is a linear functional on $MF(p)$. For the boundedness of \tilde{f} , assume ϕ is a nonzero element in $MF(p)$ and $\varphi = \frac{|\phi|}{\|\phi\|}$ then $\phi \in L^2(\varphi)$ with $\|\phi\|_{L^2(\varphi)} \leq \|\phi\|$ and

$$|\tilde{f}(\phi)| = |\langle f(\varphi), \omega_{\phi\varphi} \rangle_\varphi| \leq \|f(\varphi)\|_\varphi \|\omega_{\phi\varphi}\|_\varphi \leq \|f\|_\infty \|\phi\|_{L^2(\varphi)} \leq \|f\|_\infty \|\phi\|.$$

Consequently, $\tilde{f} \in (MF(p))^* = Mp$. When we consider \tilde{f} as an element of Mp , for any φ in $F(p)$ and x in M we have

$$\langle \tilde{f}\omega_\varphi, x\omega_\varphi \rangle_\varphi = \varphi(x^*\tilde{f}) = x^*\varphi(\tilde{f}) = \tilde{f}(x^*\varphi) = \langle f(\varphi), x\omega_\varphi \rangle_\varphi.$$

This means that the vector section $(\tilde{f}\omega_\varphi)_\varphi$ is exactly the original f .

Conversely, since the embedding $Mp \hookrightarrow \prod_{\varphi \in F(p)} H_\varphi$ is an isometry with respect to the ℓ^∞ norm, we have an isometrical isomorphism Θ from Mp onto the Banach space of all admissible vector sections f over $F(p)$ such that $\Theta(\tilde{f}) = f$. \square

We now proceed to prove the second part of theorem ??. The following easy lemma is stated for reference.

Lemma 2.9 *Let $(y_\varphi)_{\varphi \in F(p)}$ be an affine vector section over $F(p)$. If $0 \leq \lambda \leq 1$, $\varphi, \psi_1, \dots, \psi_n \in F(p)$ and $\varphi = \psi_1 + \dots + \psi_n$ then for every x in M we have*

$$\langle y_{\lambda\varphi}, x\omega_{\lambda\varphi} \rangle_{\lambda\varphi} = \lambda \langle y_\varphi, x\omega_\varphi \rangle_\varphi$$

and

$$\langle y_\varphi, x\omega_\varphi \rangle_\varphi = \langle y_{\psi_1}, x\omega_{\psi_1} \rangle_{\psi_1} + \dots + \langle y_{\psi_n}, x\omega_{\psi_n} \rangle_{\psi_n}.$$

To motivate the next step of the proof, we note that for $y\varphi$ in Mp and $0 \leq \psi \leq \varphi$ in $F(p)$ we always have, for all x in M ,

$$\psi(x^*y) = \langle y\omega_\psi, x\omega_\psi \rangle_\psi = \langle y\omega_\varphi, x\omega_{\psi\varphi} \rangle_\varphi = \langle y\omega_\varphi, T_{\psi\varphi}^*(x\omega_\psi) \rangle_\varphi = \langle T_{\psi\varphi}(y\omega_\varphi), x\omega_\psi \rangle_\psi.$$

Lemma 2.10 *Let $(y_\varphi)_{\varphi \in F(p)}$ be an affine vector section over $F(p)$. Assume φ, ψ in $F(p)$ satisfy that $0 \leq \psi \leq \varphi$. Suppose there is a projection P in the commutant $\pi_\varphi(M)'$ of $\pi_\varphi(M)$ in $B(H_\varphi)$ such that $\psi(x) = \langle x\omega_\varphi, P\omega_\varphi \rangle_\varphi, \forall x \in M$. We have*

$$\langle y_\psi, x\omega_\psi \rangle_\psi = \langle y_\varphi, x\omega_{\psi\varphi} \rangle_\varphi, \quad \forall x \in M.$$

PROOF. Write $\varphi = \psi + \rho$, where ρ in $F(p)$ is defined by

$$\rho(x) = \langle x\omega_\varphi, (1 - P)\omega_\varphi \rangle_\varphi, \quad \forall x \in M.$$

Define two isometries R from H_ψ into H_φ and S from H_ρ into H_φ by setting

$$R(x\omega_\psi) = P(x\omega_\varphi) \text{ and } S(x\omega_\rho) = (1 - P)(x\omega_\varphi), \quad \forall x \in M.$$

Note that $RH_\psi = PH_\varphi$ and $SH_\rho = (1 - P)H_\varphi$. Observe that for all x in M ,

$$\langle y_\psi, x\omega_\psi \rangle_\psi = \langle Ry_\psi, R(x\omega_\psi) \rangle_\varphi = \langle Ry_\psi, xP\omega_\varphi \rangle_\varphi$$

and

$$\langle y_\rho, x\omega_\rho \rangle_\rho = \langle Sy_\rho, S(x\omega_\rho) \rangle_\varphi = \langle Sy_\rho, x(1 - P)\omega_\varphi \rangle_\varphi.$$

By lemma ??, for every x in M

$$\begin{aligned} \langle y_\varphi, x\omega_\varphi \rangle_\varphi &= \langle y_\psi, x\omega_\psi \rangle_\psi + \langle y_\rho, x\omega_\rho \rangle_\rho \\ &= \langle Ry_\psi, xP\omega_\varphi \rangle_\varphi + \langle Sy_\rho, x(1 - P)\omega_\varphi \rangle_\varphi \\ &= \langle Ry_\psi + Sy_\rho, x\omega_\varphi \rangle_\varphi, \end{aligned}$$

since $(1 - P)Ry_\psi = PSy_\rho = 0$. Consequently, $y_\varphi = Ry_\psi + Sy_\rho$ and thus $Py_\varphi = Ry_\psi$. It is clear that $P\omega_\varphi = \omega_{\psi\varphi}$. Hence

$$\langle y_\psi, x\omega_\psi \rangle_\psi = \langle Ry_\psi, R(x\omega_\psi) \rangle_\varphi = \langle Py_\varphi, xP\omega_\varphi \rangle_\varphi = \langle y_\varphi, x\omega_{\psi\varphi} \rangle_\varphi.$$

□

PROOF OF THE SECOND PART OF THEOREM ??. Let $(y_\varphi)_{\varphi \in F(p)}$ be a bounded affine vector section over $F(p)$. We prove that for every φ and ψ in $F(p)$ such that $0 \leq \psi \leq \varphi$,

$$\langle y_\psi, x\omega_\psi \rangle_\psi = \langle y_\varphi, x\omega_{\psi\varphi} \rangle_\varphi, \quad \forall x \in M.$$

By the Radon–Nikodym theorem (see *e.g.* Sakai [?]), there is a T in $\pi_\varphi(M)'$, $0 \leq T \leq 1$, such that $\psi(x) = \langle x\omega_\varphi, T\omega_\varphi \rangle_\varphi, \forall x \in M$, *i.e.* $T\omega_\varphi = \omega_{\psi\varphi}$. By the spectral theorem for bounded self-adjoint Hilbert space operators, we can write

$$T = \int_0^1 \lambda dE(\lambda),$$

where E is the projection-valued measure related to T . For $\varepsilon > 0$, there is a partition $\{\Delta_1, \dots, \Delta_n\}$ of $[0, 1]$ and $\lambda_1, \dots, \lambda_n$ between 0 and 1 such that $0 \leq \sum \lambda_k E(\Delta_k) \leq T$ and $\|T - \sum \lambda_k E(\Delta_k)\| < \varepsilon$. Define ψ_k in $F(p)$ by

$$\psi_k(x) = \langle x\omega_\varphi, E(\Delta_k)\omega_\varphi \rangle_\varphi, \quad \forall x \in M, k = 1, \dots, n.$$

It is equivalent to say that

$$E(\Delta_k)\omega_\varphi = \omega_{\psi_k\varphi}, \quad k = 1, \dots, n.$$

Since $E(\Delta_k) \in \pi_\varphi(M)'$, $k = 1, \dots, n$, we have, by lemma ??,

$$\langle y_{\psi_k}, x\omega_{\psi_k} \rangle_{\psi_k} = \langle y_\varphi, x\omega_{\psi_k\varphi} \rangle_\varphi = \langle y_\varphi, xE(\Delta_k)\omega_\varphi \rangle_\varphi.$$

Let $\psi_0 = \sum \lambda_k \psi_k$. We have $0 \leq \psi_0 \leq \psi \leq \varphi$. Write $\psi = \psi_0 + \rho$. Note $\rho \in F(p)$ and

$$\|\rho\| = \|\psi\| - \|\psi_0\| = \left\langle \omega_\varphi, (T - \sum \lambda_k E(\Delta_k))\omega_\varphi \right\rangle_\varphi \leq \|T - \sum \lambda_k E(\Delta_k)\| \|\varphi\| < \|\varphi\| \varepsilon.$$

By lemma ??,

$$\begin{aligned} \langle y_\psi, x\omega_\psi \rangle_\psi &= \langle y_{\psi_0}, x\omega_{\psi_0} \rangle_{\psi_0} + \langle y_\rho, x\omega_\rho \rangle_\rho \\ &= \sum_{k=1}^n \lambda_k \langle y_{\psi_k}, x\omega_{\psi_k} \rangle_{\psi_k} + \langle y_\rho, x\omega_\rho \rangle_\rho \\ &= \sum_{k=1}^n \lambda_k \langle y_\varphi, xE(\Delta_k)\omega_\varphi \rangle_\varphi + \langle y_\rho, x\omega_\rho \rangle_\rho \\ &= \left\langle y_\varphi, x \sum_{k=1}^n \lambda_k E(\Delta_k)\omega_\varphi \right\rangle_\varphi + \langle y_\rho, x\omega_\rho \rangle_\rho. \end{aligned}$$

Therefore,

$$\begin{aligned} & \left| \langle y_\psi, x\omega_\psi \rangle_\psi - \langle y_\varphi, x\omega_{\psi\varphi} \rangle_\varphi \right| \\ & \leq \left| \left\langle y_\varphi, x \left(T - \sum_{k=1}^n \lambda_k E(\Delta_k) \right) \omega_\varphi \right\rangle_\varphi \right| + \left| \langle y_\rho, x\omega_\rho \rangle_\rho \right| \\ & \leq \|y_\varphi\|_\varphi \|x\| \|T - \sum_{k=1}^n \lambda_k E(\Delta_k)\| \|\omega_\varphi\|_\varphi + \|y_\rho\|_\rho \|x\| \|\omega_\rho\|_\rho \\ & < K \|x\| \|\varphi\|^{1/2} \varepsilon + K \|x\| \|\varphi\|^{1/2} \varepsilon^{1/2}, \end{aligned}$$

where $K = \|y\|_\infty$ is the bound of y . Since ε is arbitrary, $\langle y_\psi, x\omega_\psi \rangle_\psi = \langle y_\varphi, x\omega_{\psi\varphi} \rangle_\varphi = \langle y_\varphi, T_{\psi\varphi}^*(x\omega_\psi) \rangle_\varphi$, $\forall x \in M$. In other words, $T_{\psi\varphi} y_\varphi = y_\psi$, as asserted. Thus $(y_\varphi)_\varphi$ is admissible. The fact that every admissible vector section is bounded and affine follows from the first part of the proof. \square

3 Represent C*-algebras via continuous admissible vector sections

Let A be a C*-algebra and p the closed projection in A^{**} related to a closed left ideal L of A . Let $F(p) = \{\varphi \in A^* : \varphi \geq 0, \varphi(p) = \|\varphi\| \leq 1\}$. As a special case of theorem ??, we have

Theorem 3.1 $A^{**}p$ ($\cong A^{**}/L^{**}$) is isometrically isomorphic to the Banach space of all admissible vector sections over $F(p)$, which consists exactly of all bounded affine vector sections over $F(p)$.

It is natural to ask which admissible vector sections Ap contains. Analogous to the classical Kadison function representation (cf. [?]) one may guess Ap consists of all “continuous” affine vector sections over $F(p)$. The question is how we define continuity for the field $(F(p), \{H_\varphi\}_\varphi)$ of Hilbert spaces. Of course, all vector sections arising from Ap should be continuous.

Recall the notion of a continuous field of (complex) Hilbert spaces [?, ?]. Let T be a Hausdorff space called the *base space*. For each t in T , let H_t be a (complex) Hilbert space, called the *fiber Hilbert space*. A *vector section* is a function x on T such that $x(t) \in H_t, \forall t \in T$. A (*full*) *continuous structure* for the field $(T, \{H_t\}_{t \in T})$ of Hilbert spaces is a linear space Γ of vector sections, called *continuous vector sections*, satisfying the conditions:

- (i) $t \mapsto \|x(t)\|_{H_t}$ is continuous on T for all x in Γ .
- (ii) $\{x(t) : x \in \Gamma\}$ is norm dense in H_t for all t in T .
- (iii) Let x be a vector section; if for any t in T and $\varepsilon > 0$ there exists an a in Γ such that $\|x(t) - a(t)\| < \varepsilon$ throughout a neighborhood of t then $x \in \Gamma$.

The triple $(T, \{H_t\}_t, \Gamma)$ is called a *continuous field of Hilbert spaces*.

A linear space X of vector sections which satisfies conditions (i) and (ii) defines a continuous structure $\Gamma(X)$, which is the set of all vector sections x satisfying the condition that

- (iii)' For any t in T and $\varepsilon > 0$ there exists an a in X such that $\|x(t) - a(t)\| < \varepsilon$ throughout a neighborhood of t .

It is easy to see that for a vector section $x = (x(t))_{t \in T}$, x is continuous (that is $x \in \Gamma(X)$) if and only if

1. $t \mapsto \langle x(t), x(t) \rangle_{H_t}$ is continuous on T , and
2. $t \mapsto \langle x(t), y(t) \rangle_{H_t}$ is continuous on T , $\forall y \in X$.

A vector section x is said to be *bounded* if $\|x\|_\infty = \sup_{t \in T} \|x(t)\|_{H_t} < \infty$. x is said to be *weakly continuous* in $(T, \{H_t\}_t, \Gamma(X))$ if the scalar function $t \mapsto \langle x(t), y(t) \rangle_{H_t}$ is continuous on T for every y in $\Gamma(X)$. If x is bounded then $\Gamma(X)$ can be replaced by X in the above condition. A weakly continuous vector section x is continuous if and only if $t \mapsto \langle x(t), x(t) \rangle_{H_t}$ is continuous on T (cf. [?]).

Now, let us point out that if $ap \in Ap$ then $\varphi \mapsto \|a\omega_\varphi\|_\varphi$ is continuous on $F(p)$. It is also clear that $A\omega_\varphi$ is norm dense in H_φ for each φ in $F(p)$. Therefore, the set X of vector sections arising from Ap defines a continuous structure $\Gamma(X)$ which we shall henceforth write as $\Gamma(Ap)$. A vector section $(x_\varphi)_{\varphi \in F(p)}$ in $(F(p), \{H_\varphi\}_\varphi, \Gamma(Ap))$ is continuous if and only if for any $\varepsilon > 0$ and φ in $F(p)$ there exist an a in A and a neighborhood V_φ of φ in $F(p)$ such that

$$\|x_\psi - a\omega_\psi\|_\psi < \varepsilon, \quad \forall \psi \in V_\varphi.$$

In this context, a bounded vector section $(x_\varphi)_{\varphi \in F(p)}$ is *weakly continuous* if $\varphi \mapsto \langle x_\varphi, a\omega_\varphi \rangle_\varphi$ is continuous on $F(p)$, $\forall a \in A$. A weakly continuous vector section $(x_\varphi)_{\varphi \in F(p)}$ is *continuous* if $\varphi \mapsto \langle x_\varphi, x_\varphi \rangle_\varphi$ is continuous on $F(p)$. Moreover, a vector section $(x_\varphi)_{\varphi \in F(p)}$ is continuous if and only if $\varphi \mapsto \langle x_\varphi, y_\varphi \rangle_\varphi$ is continuous on $F(p)$ for all weakly continuous vector sections $(y_\varphi)_{\varphi \in F(p)}$. In fact, $(x_\varphi)_{\varphi \in F(p)}$ itself must be weakly continuous in this case, and thus $\varphi \mapsto \langle x_\varphi, x_\varphi \rangle_\varphi$ is continuous on $F(p)$, too. It is plain that continuous vector sections need not arise from elements of Ap . However, we have

Theorem 3.2 *$Ap (\cong A/L)$ is isometrically isomorphic to the Banach space of all continuous admissible (= continuous and affine) vector sections of the continuous field of Hilbert spaces $(F(p), \{H_\varphi\}_\varphi, \Gamma(A))$.*

PROOF. We adopt the notations used in the last section with M replaced by A^{**} . Let $f = (f(\varphi))_\varphi$ be a continuous admissible vector section over $F(p)$. In view of theorem ??, it suffices to show that whenever $\phi_\lambda \rightarrow \phi$ in the weak* topology of the polar $L^\circ = (A/L)^*$ of L in A^* , $\tilde{f}(\phi_\lambda) \rightarrow \tilde{f}(\phi)$. By the Krein–Smulian theorem, we need only to check this for bounded nets. So assume $\|\phi_\lambda\| \leq 1$. Note that $L^\circ = \{\psi \in A^* : \psi = \psi(\cdot p)\}$

and hence if $\psi \in L^\circ$ and $\|\psi\| \leq 1$ then $|\psi| \in F(p)$. Since $F(p)$ is weak* compact, there is a subnet ϕ_k of ϕ_λ such that $\varphi_k = |\phi_k|$ converges to an element φ of $F(p)$ in the weak* topology (note that φ is not necessarily $|\phi|$, see *e.g.* [?]). Now for any a in A the inequalities

$$|\phi_k(a)|^2 \leq \|\varphi_k\| \varphi_k(a^*a), \quad \forall k,$$

imply

$$|\phi(a)|^2 \leq K\varphi(a^*a).$$

Here $K = \sup_k \|\varphi_k\| \leq 1$. Therefore, ϕ is observable at φ and thus

$$\tilde{f}(\phi) = \langle f(\varphi), \omega_{\phi\varphi} \rangle_\varphi.$$

Let $\varepsilon > 0$. Since f is a continuous vector section in $(F(p), \{H_\varphi\}_\varphi, \Gamma(Ap))$, there exist a neighborhood U_φ of φ in $F(p)$ and an a in A such that $\|f(\psi) - a\omega_\psi\|_\psi < \varepsilon/3$ in U_φ . Thus

$$\|f(\varphi) - a\omega_\varphi\|_\varphi < \varepsilon/3$$

and

$$\|f(\varphi_k) - a\omega_{\varphi_k}\|_{\varphi_k} < \varepsilon/3,$$

eventually. Also,

$$|\phi(a) - \phi_k(a)| < \varepsilon/3$$

eventually. So for k sufficiently large,

$$\begin{aligned} & \left| \tilde{f}(\phi) - \tilde{f}(\phi_k) \right| \\ &= \left| \langle f(\varphi), \omega_{\phi\varphi} \rangle_\varphi - \langle f(\varphi_k), \omega_{\phi_k\varphi_k} \rangle_{\varphi_k} \right| \\ &\leq \left| \langle f(\varphi), \omega_{\phi\varphi} \rangle_\varphi - \langle a\omega_\varphi, \omega_{\phi\varphi} \rangle_\varphi \right| + \left| \langle a\omega_\varphi, \omega_{\phi\varphi} \rangle_\varphi - \langle a\omega_{\varphi_k}, \omega_{\phi_k\varphi_k} \rangle_{\varphi_k} \right| \\ &\quad + \left| \langle a\omega_{\varphi_k}, \omega_{\phi_k\varphi_k} \rangle_{\varphi_k} - \langle f(\varphi_k), \omega_{\phi_k\varphi_k} \rangle_{\varphi_k} \right| \\ &\leq \|f(\varphi) - a\omega_\varphi\|_\varphi + |\phi(a) - \phi_k(a)| + \|a\omega_{\varphi_k} - f(\varphi_k)\|_{\varphi_k} \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon. \end{aligned}$$

Consequently, $\tilde{f}(\phi_k) \longrightarrow \tilde{f}(\phi)$. Since the same argument can be applied to any subnet of ϕ_λ , we have $\tilde{f}(\phi_\lambda) \longrightarrow \tilde{f}(\phi)$. Hence f defines an element in Ap , as asserted. \square

4 Continuity and weak continuity

From now on, a continuous admissible (resp. admissible) vector section is considered as an element of Ap (resp. $A^{**}p$). Denote by \mathcal{W}_p the family of all weakly continuous admissible vector sections over $F(p)$.

Corollary 4.1 *Let $xp \in A^{**}p$.*

1. $px^*xp \in pAp$ and $pa^*xp \in pAp, \quad \forall ap \in Ap \Leftrightarrow xp \in Ap$.
2. $pa^*xp \in pAp, \forall ap \in Ap \Leftrightarrow xp \in \mathcal{W}_p$.
3. $pw^*xp \in pAp, \forall wp \in \mathcal{W}_p \Leftrightarrow xp \in Ap$.

PROOF. It follows from [?, 3.5] that for x, y in A^{**} , $\varphi \mapsto \langle x\omega_\varphi, y\omega_\varphi \rangle_\varphi = \varphi(y^*x)$ is continuous on $F(p)$ if and only if $py^*xp \in pAp$. Recalling the discussion of fields of Hilbert spaces in section 3, we see that (??) and (??) are immediate whilst (??) is just a restatement of theorem ??.

In case $p = 1$, an admissible vector section xp is weakly continuous if and only if $x \in RM(A)$, the set of right multipliers of A (cf. [?]). In general, we have $RM(A)p \subseteq \mathcal{W}_p$. To investigate what \mathcal{W}_p contains, we quote a result of Brown [?, 3.9]:

Theorem 4.2 *Let A be a σ -unital C^* -algebra and p a closed projection in A^{**} . Let xp in $A^{**}p$ be such that $\|xp\| = 1$ and $Axp \subseteq Ap$. Then there is a right multiplier r of A in A^{**} such that $\|r\| = 1$ and $xp = rp$.*

Corollary 4.3 *If A is a σ -unital C^* -algebra and p is a closed, central projection in A^{**} then $\mathcal{W}_p = RM(A)p$.*

Corollary 4.4 *Let A be a σ -unital C^* -algebra and p a closed projection in A^{**} . For an xp in \mathcal{W}_p ,*

$$xp \in RM(A)p \Leftrightarrow px^*Axp \subseteq pAp.$$

PROOF. One direction is obvious. For the other one, we assume $xp \notin RM(A)p$. Then there is an a in A such that $axp \notin Ap$ by theorem ??. Since axp is also a weakly continuous vector section, we must have $px^*a^*axp \notin pAp$. Hence, px^*Axp is not contained in pAp .

Corollary 4.5 *Let A be a σ -unital C^* -algebra and p a closed projection in A^{**} .*

1. *If $xp \in \mathcal{W}_p$ and $xp = pxp$ then $xp \in RM(A)p$.*
2. *If A is simple and $p \in M(A)$ then $\mathcal{W}_p = RM(A)p$.*

PROOF. (??) We check the condition $px^*Axp \subseteq pAp$. In fact,

$$\begin{aligned} px^*Axp &= px^*Apxp \subseteq pApxp, \text{ since } xp \in \mathcal{W}_p, \\ &= pAxp \subseteq pAp, \text{ again since } xp \in \mathcal{W}_p. \end{aligned}$$

(??) Since ApA is an ideal of A and A is simple, either $ApA = \{0\}$ or the norm closure \overline{ApA} of ApA coincides with A . But $ApA = \{0\}$ implies $p = 0$. The assertion becomes trivial in this case. So assume $\overline{ApA} = A$. Now if $xp \in \mathcal{W}_p$, we have

$$px^*Axp = px^*\overline{ApA}xp \subseteq \overline{pApAp} \subseteq pAp.$$

The proof is complete since it is always true that $RM(A)p \subseteq \mathcal{W}_p$. □

In the following we present an example to show that the conclusions of corollary ?? can fail if the hypothesis in (??) or (??) is not fulfilled.

Example 4.6 Let H be a separable infinite dimensional Hilbert space with an orthonormal basis $\{e_1, e_2, \dots\}$. Let p be the projection of H onto $\text{span}\{e_1, e_3, e_5, \dots\}$ and $A = C^*(\mathcal{K}, 1 - p)$, the C^* -subalgebra of $B(H)$ generated by \mathcal{K} , the C^* -subalgebra of all compact operators on H , and $1 - p$. Then the separable (hence σ -unital) C^* -algebra A is given by

$$A = \{T + \lambda(1 - p) : T \in \mathcal{K}, \lambda \in \mathbb{C}\}$$

and A^{**} can be described as

$$A^{**} = B(H) \oplus \mathbb{C}(1 - p).$$

When A^{**} is viewed in this way, the embedding of A into A^{**} is given by

$$T + \lambda(1 - p) \longrightarrow (T + \lambda(1 - p), \lambda(1 - p)).$$

Identify p with $p \oplus 0$ in A^{**} . Then $p \in M(A)$. Note that A is not simple and thus corollary ??(??) does not apply. It is easy to see that $Ap = \mathcal{K}p$, $pAp = p\mathcal{K}p$ and $\mathcal{W}_p = B(H)p$. On the other hand,

$$RM(A) = \{(K + \lambda(1 - p) + pS, \lambda(1 - p)) : K \in \mathcal{K}, S \in B(H) \text{ and } \lambda \in \mathbb{C}\}.$$

Hence $RM(A)p = \mathcal{K}p + pB(H)p$. It is clear that $\mathcal{W}_p \neq RM(A)p$. For example, if T is the unilateral shift, *i.e.* $Te_n = e_{n+1}, n = 1, 2, \dots$ then $Tp \in \mathcal{W}_p$ but $Tp \notin RM(A)p$ (since $(1-p)Tp = Tp \notin Ap$). We also note that $Tp \neq pTp = 0$ and thus corollary ??(??) does not apply, either. \square

5 Comparison with Takesaki duality theorem

Let A be a C^* -algebra. Let H be a Hilbert space of sufficiently large infinite dimension such that every cyclic representation of A is unitarily equivalent to a cyclic representation of A on H . Let p_π be the projection of H onto $H_\pi = \overline{\pi(A)H}^{\|\cdot\|}$ for each π in the set $Rep(A, H)$ of all representations of A on H . For each partial isometry u in $B(H)$ and π in $Rep(A, H)$ such that $u^*u \geq p_\pi$, we denote by π^u the representation $u\pi u^*$, *i.e.* $\pi^u(a) = u\pi(a)u^*, \forall a \in A$. We equip $Rep(A, H)$ the point strong operator topology (PSOT):

$$\pi_\lambda \xrightarrow{\text{PSOT}} \pi \text{ in } Rep(A, H) \text{ if } \pi_\lambda(a)h \xrightarrow{\|\cdot\|} \pi(a)h \text{ in } H, \quad \forall a \in A, \forall h \in H.$$

Definition 5.1 ([?], [?]) A function $T : Rep(A, H) \longrightarrow B(H)$ is said to be a *TB-admissible operator field* if the following conditions are satisfied:

$$(TB_1) \quad \|T\| := \sup\{\|T(\pi)\| : \pi \in Rep(A, H)\} < \infty.$$

$$(TB_2) \quad T(\pi) = p_\pi T(\pi) = T(\pi)p_\pi, \forall \pi \in Rep(A, H).$$

$$(TB_3) \quad T(\pi + \pi') = T(\pi) + T(\pi') \text{ whenever } \pi, \pi' \in Rep(A, H) \text{ such that } H_\pi \perp H_{\pi'}.$$

$$(TB_4) \quad T(\pi_u) = uT(\pi)u^* \text{ whenever } \pi \in Rep(A, H) \text{ and } u \text{ is a partial isometry in } B(H) \text{ such that } u^*u \geq p_\pi.$$

In [?], Bichteler extended Takesaki duality theorem [?] for separable C^* -algebras A to the general form:

Theorem 5.2 *The set of all TB-admissible operator fields is isometrically isomorphic to A^{**} in the sense that for each TB-admissible operator field $T = (T(\pi))_\pi$ there is a t in A^{**} such that*

$$\pi(t) = T(\pi), \quad \forall \pi \in Rep(A, H).$$

(Here π is understood to be (uniquely) extended to a $\sigma(A^{**}, A^*)$ -continuous representation (again denoted by π) of A^{**} on H .) Moreover, $t \in A$ if and only if T is PSOT-SOT

continuous in the sense that if $\pi_\lambda \xrightarrow{\text{PSOT}} \pi$ in $\text{Rep}(A, H)$ then $T(\pi_\lambda) \longrightarrow T(\pi)$ in $B(H)$ with the strong operator topology (SOT).

A similar argument as in the proof of proposition ?? gives

Proposition 5.3 *Every function $T : \text{Rep}(A, H) \longrightarrow B(H)$ which satisfies (TB_2) , (TB_3) and (TB_4) is TB-admissible. In other words, (TB_1) is redundant.*

Definition 5.4 Let $\pi \in \text{Rep}(A, H)$ and $h \in H$ with $\|h\| \leq 1$. Let φ in $Q(A)$ be defined by $\varphi := \langle \pi(\cdot)h, h \rangle_H$. We define an isometry $U_{\pi, h}^\varphi$ from H_φ into H by

$$U_{\pi, h}^\varphi(a\omega_\varphi) := \pi(a)h, \quad \forall a \in A,$$

where $a\omega_\varphi$ denotes $\pi_\varphi(a)\omega_\varphi$ in the GNS representation $(\pi_\varphi, H_\varphi, \omega_\varphi)$ induced by φ , as before.

Some lengthy computation and straightforward reasoning will bring us the following connection of Takesaki duality theorem and our representation theory developed in earlier sections in this paper.

Theorem 5.5 ([?]) *There exists an isometrical isomorphism from the Banach space of all admissible vector sections $x = (x_\varphi)_\varphi$ over $Q(A)$ onto the Banach space of all TB-admissible operator fields $T = (T(\pi))_\pi$ such that the relation*

$$U_{\pi, h}^\varphi x_\varphi = T(\pi)h$$

is satisfied whenever $\varphi = \langle \pi(\cdot)h, h \rangle_H$ for some π in $\text{Rep}(A, H)$ and h in H with $\|h\| \leq 1$. Moreover, $T = (T(\pi))_\pi$ is a continuous TB-admissible operator field if and only if $x = (x_\varphi)_\varphi$ is a continuous admissible vector section.

Roughly speaking, Takesaki [?] represented x in A^{**} as a field of operators (matrices) $\pi(x)$'s and we represent x as a field of vectors (columns) $x\omega_\varphi$'s. The general version of our representation of $A^{**}p$ is to pay attention only on those columns $x\omega_\varphi$'s of the matrix $\pi(x)$ in the range of the closed projection p (i.e. φ is supported by p , or equivalently, $p\omega_\varphi = \omega_\varphi$). Moreover, xp comes from Ap if and only if xp has continuous coordinates $\varphi \longmapsto \langle x\omega_\varphi, a\omega_\varphi \rangle_\varphi, \forall a \in A$, and continuous norm $\varphi \longmapsto \langle x\omega_\varphi, x\omega_\varphi \rangle_\varphi^{1/2}$ over $F(p)$. In this sense, our results extend Takesaki duality theorem.

References

- [1] C. A. Akemann, “The general Stone-Weierstrass problem”, *J. Funct. Anal.*, vol. 4, 1969, 277–294.
- [2] C. A. Akemann, “Left ideal structure of C^* -algebras”, *J. Funct. Anal.*, vol. 6, 1970, 305–317.
- [3] C. A. Akemann, “A Gelfand representation theory for C^* -algebras”, *Pacific J. Math.*, vol. 39, 1971, 1-11.
- [4] C. A. Akemann, G. K. Pedersen and J. Tomiyama, “Multipliers of C^* -algebras”, *J. Funct. Anal.*, vol. 13, 1973, 277-301.
- [5] C. A. Akemann and F. Shultz, “Perfect C^* -algebras”, *Memoirs A. M. S.*, vol. 326, 1985.
- [6] K. Bichteler, “A generalization to the non-separable case of Takesaki’s duality theorem for C^* -algebras”, *Inventiones Math.*, vol. 9, 1969, 89–98.
- [7] L. G. Brown, “Semicontinuity and multipliers of C^* -algebras”, *Can. J. Math.*, vol. XL, 4, 1988, 865–988.
- [8] L. G. Brown, “Complements to various Stone-Weierstrass theorems for C^* -algebras and a theorem of Shultz”, *Commun. Math. Phys.*, vol. 143, 1992, 405–413.
- [9] J. Dixmier and A. Douady, “Champs continus d’espaces hilbertiens et de C^* -algèbres”, *Bull. Soc. Math. France*, vol. 91, 1963, 227–284.
- [10] J. Dixmier, “ C^* -algebras”, North-Holland publishing company, Amsterdam–New York–Oxford, 1977.
- [11] E. G. Effros, “Order ideals in C^* -algebras and its dual”, *Duke Math.*, vol. 30, 1963, 391–412.
- [12] J.M.G. Fell, “The structure of algebras of operator fields”, *Acta Math.*, vol. 106, 1961, 233–280.
- [13] G. K. Pedersen, “ C^* -algebras and their automorphism groups”, Academic Press, 1979, London.

- [14] R. T. Prosser “On the ideal structure of operator algebras”, *Memoirs A. M. S.*, vol. 45, 1963.
- [15] M. A. Rieffel, “Induced Representations of C^* -algebras”, *Adv. in Math.*, vol. 13, 1974, 176–257.
- [16] S. Sakai, “ C^* -algebras and W^* -algebras”, Springer–Verlag, New York, 1971.
- [17] F. W. Shultz, “Pure states as a dual object for C^* -algebra”, *Commun. Math. Phys.*, vol. 82, 1982, 497–509.
- [18] M. Takesaki, “A duality in the representation theory of C^* -algebras”, *Ann. Math.*, vol. 85, 1967, 370–382.
- [19] M. Tomita, “Spectral theory of operator algebras, I”, *Math. J. Okayama Univ.*, 1959, 63–98.
- [20] M. Tomita, “Spectral theory of operator algebras, II”, *Math. J. Okayama Univ.*, 1960, 19–60.
- [21] Ngai-ching Wong, “The left quotient of a C^* -algebra and its representation through a continuous field of Hilbert spaces”, Ph.D. Thesis, Purdue University, West Lafayette, U.S.A., 1991.
- [22] Ngai-ching Wong and L. G. Brown, “Left quotients of a C^* -algebra, II: Continuity on extreme boundary”, *preprint*.
- [23] Ngai-ching Wong and L. G. Brown, “Left quotients of a C^* -algebra, III: Operators on left quotients”, *preprint*.