

國立中山大學

NATIONAL SUN YAT-SEN UNIVERSITY

線性代數 (一)

MATH 103 / GEAI 1215: Linear Algebra I

期末考

January 7, 2019

Final Examination

姓名 Name : _____

學號 Student ID # : _____

Lecturer: Jephian Lin 林晉宏

Contents: cover page,
9 pages of questions,
score page at the end

To be answered: on the test paper

Duration: **110 minutes**

Total points: **35 points** + 7 extra points

Do not open this packet until instructed to do so.

Instructions:

- Enter your **Name** and **Student ID #** before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining it or circling it**. If multiple answers are shown then no marks will be awarded.
- 可用中文或英文作答

1. [5pt] Find the general solution of the following linear system.

$$\begin{cases} w - 2x + 3y - z = 2 \\ 2w - 4x + 6y - 1z = 6 \\ 3w - 6x + 9y - 3z = 6 \end{cases}$$

That is, find \vec{p} and $\vec{\beta}_1, \dots, \vec{\beta}_k$ such that

$$\{\vec{p} + c_1\vec{\beta}_1 + \dots + c_k\vec{\beta}_k : c_1, \dots, c_k \in \mathbb{R}\}$$

is the set of all solutions.

2. [5pt] Suppose $S = \{\vec{w}_1, \dots, \vec{w}_k\}$ is a set of nonzero vectors in \mathbb{R}^n such that $\vec{w}_i \cdot \vec{w}_j = 0$ for any distinct i and j . (That is, any two vectors in S are orthogonal to each other.) Show that S is linearly independent.

3. [3pt] Let $\vec{x} = \begin{bmatrix} 3+i \\ i \\ 2+i \end{bmatrix}$ and $\vec{y} = \begin{bmatrix} 1-2i \\ 2+3i \\ 1 \end{bmatrix}$. Find the values of the inner products $\langle \vec{x}, \vec{y} \rangle$, $\langle \vec{y}, \vec{x} \rangle$ and the norm $|\vec{x}|$ in \mathbb{C}^3 .

4. [2pt] Let $\vec{u} = \begin{bmatrix} 4 \\ 2 \\ 3 \end{bmatrix}$ and $\vec{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$. Find \vec{x} and \vec{y} such that $\vec{u} = \vec{x} + \vec{y}$ with $\vec{x} \in \text{span}\{\vec{v}\}$ and $\langle \vec{v}, \vec{y} \rangle = 0$.

5. [5pt] Let $V = \text{span} \left\{ \begin{bmatrix} 1 \\ -2 \\ 3 \\ -1 \end{bmatrix}, \begin{bmatrix} 2 \\ -4 \\ 6 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ -6 \\ 9 \\ -3 \end{bmatrix} \right\}$. Find a basis of V and a basis of V^\perp .

6. [5pt] Consider the following data:

x	1	2	3	4	5
y_i	1	2	2	3	3

Find a line $f(x) = ax + b$ such that the error

$$\sum_{i=1}^5 (f(x_i) - y_i)^2$$

is minimized.

[The orthogonal projection of a vector \vec{v} onto the column space of a matrix \mathbf{A} is $\mathbf{A}(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \vec{v}$. Your answer can be a formula without computing the final answer, but you have to specify all matrices or vectors occurred in your formula.]

7. [5pt] Let $f : V \rightarrow W$ be a homomorphism. Show that $f^{-1}(Y)$ is a subspace of V if Y is a subspace of W .

8. [5pt] Let $f : V \rightarrow W$ be a homomorphism. Show that f is one-to-one if and only if $\text{nullspace}(f) = \{\vec{\mathbf{0}}\}$.

9. Let

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 10 \\ 1 \\ 0 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 5 \\ 10 \\ 1 \end{bmatrix}, \text{ and } \vec{u} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}.$$

Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a homomorphism such that

$$f(\vec{v}_1) = f(\vec{v}_2) = f(\vec{v}_3) = \vec{u}.$$

(a) [extra 1pt] Find $\text{range}(f)$ in set notation and give a brief reason to your answer. [Hint: $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ is a basis of \mathbb{R}^3 .]

(b) [extra 2pt] Find the rank of f and the nullity of f .

(c) [extra 2pt] Find a basis of $\text{nullspace}(f)$ and give a brief reason to your answer.

10. [extra 2pt] Recall that the trace of a matrix is the sum of its diagonal entries. That is,

$$\operatorname{tr}(A) = \sum_{i=1}^n a_{i,i}$$

if $A = [a_{i,j}]$ is an $n \times n$ matrix. Let $\mathcal{M}_{n \times n}$ be the set of all $n \times n$ matrices. Then

$$\begin{aligned} f : \mathcal{M}_{n \times n} &\rightarrow \mathbb{R} \\ A &\mapsto \operatorname{tr}(A) \end{aligned}$$

is a homomorphism. (You don't have to check this fact.) Find the rank of f and the nullity of f in terms of n .

[END]

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	5	
6	5	
7	5	
8	5	
9	2	
Total	35 (+7)	