

# LEFT QUOTIENTS OF C\*-ALGEBRAS, II: ATOMIC PARTS OF LEFT QUOTIENTS

NGAI-CHING WONG

ABSTRACT. Let  $A$  be a C\*-algebra. Let  $z$  be the maximal atomic projection in  $A^{**}$ . By a theorem of Brown,  $x$  in  $A^{**}$  has a continuous atomic part, *i.e.*  $zx = za$  for some  $a$  in  $A$ , whenever  $x$  is uniformly continuous on the set of pure states of  $A$ . Let  $L$  be a closed left ideal of  $A$ . Under some additional conditions, we shall show that for any  $x$  in  $A^{**}$ ,  $x$  has a continuous atomic part modulo  $L^{**}$ , *i.e.*  $zx + L^{**} = za + L^{**}$  for some  $a$  in  $A$  whenever  $x^*x$  and  $u^*x$ ,  $\forall u \in A$ , are uniformly continuous on the set of pure states of  $A$  vanishing on  $L$ .

## 1. INTRODUCTION

Let  $A$  be a C\*-algebra with the Banach dual  $A^*$  and double dual  $A^{**}$ . Let

$$Q(A) = \{\varphi \in A^* : \varphi \geq 0, \|\varphi\| \leq 1\}$$

be the quasi-state space of  $A$ .  $Q(A)$  is a weak\* compact convex set. The extreme boundary of  $Q(A)$  is the pure state space  $P(A) \cup \{0\}$  of  $A$ . In the Kadison function representation, elements  $x$  of  $A^{**}$  are represented as bounded and affine functionals  $\varphi \mapsto \varphi(x)$  of  $Q(A)$  vanishing at zero.  $x \in A$  if and only if  $x$  is continuous on  $Q(A)$  (see, *e.g.* [16]).

Let  $z$  be the maximal atomic projection in  $A^{**}$ . Note that  $A^{**} = zA^{**} \oplus (1-z)A^{**}$ ;  $zA^{**}$  is the direct sum of type I factors and  $(1-z)A^{**}$  has no type I factor direct summand of  $A^{**}$ . In particular,  $z$  is a central projection in  $A^{**}$  supporting all pure states of  $A$ . In other words,  $\varphi(x) = \varphi(zx)$  for all  $x$  in  $A^{**}$  and all pure states  $\varphi$  of  $A$ .

$zA^{**}$  is called the *atomic part* of  $A^{**}$ . An element  $x$  of  $A^{**}$  is said to *have a continuous atomic part* if  $zx = za$  for some  $a$  in  $A$  (cf. [18]). In this case,  $x$  and  $a$  agree on  $P(A) \cup \{0\}$  since  $\varphi(x) = \varphi(zx) = \varphi(za) = \varphi(a)$  for all pure states  $\varphi$  of  $A$ . In particular,  $\varphi \mapsto \varphi(x)$  is uniformly continuous on  $P(A) \cup \{0\}$ . The converse is also true. Shultz [18] showed that  $x$  in  $A^{**}$  has a continuous atomic part whenever  $x, x^*x$  and  $xx^*$  are uniformly continuous on  $P(A) \cup \{0\}$ . Recently, Brown [7] got the complete result.

**Theorem 1.1** ([7]). *Let  $x \in A^{**}$ .  $x$  has a continuous atomic part (i.e.  $zx \in zA$ ) if and only if  $x$  is uniformly continuous on  $P(A) \cup \{0\}$ .*

---

*Date:* May 27, 1998; a version submitted to J. Operator Theory.

*1991 Mathematics Subject Classification.* 46L05, 46L45.

*Key words and phrases.* C\*-algebras, continuous fields of Hilbert spaces, continuous atomic parts.

Partially supported by National Science Council of Republic of China, NSC 86-2115-M110-002.

In view of the Kadison function representation, Theorem 1.1 merely states that a uniformly continuous functional of the extreme boundary  $P(A) \cup \{0\}$  of  $Q(A)$  can be lifted to a continuous affine functional of the whole of  $Q(A)$ .

Let  $L$  be a norm closed left ideal of the  $C^*$ -algebra  $A$ . The Banach double dual  $L^{**}$  of  $L$  is a weak\* closed left ideal of the  $w^*$ -algebra  $A^{**}$ .

**Definition 1.2.** Let  $x \in A^{**}$ .  $x$  is said to *have a continuous atomic part modulo  $L^{**}$*  if  $zx + L^{**} = za + L^{**}$  for some  $a$  in  $A$ .

We are interested in the question when  $x$  in  $A^{**}$  has a continuous atomic part modulo  $L^{**}$ . By applying our tools developed in this paper to the case  $L = \{0\}$ , we shall get new and interesting results. As a corollary of Theorems 1.1 and 1.7 below, for example, we have

**Theorem 1.3.** *Let  $x \in A^{**}$ .  $x$  has a continuous atomic part (i.e.  $zx \in zA$ ) if and only if  $x^*x$  and  $a^*x$ ,  $\forall a \in A$ , are uniformly continuous on  $P(A) \cup \{0\}$ .*

Theorem 1.3 supplements results of Shultz [18] and Brown [7] in the case  $A$  is non-unital. More general statements appear in succeeding sections.

We are going to give a precise meaning of our assertions. Recall that a convex subset  $F$  of a convex set  $Q$  is called a *face* of  $Q$  if we can always infer that  $\varphi$  and  $\psi$  belong to  $F$  whenever  $\varphi, \psi \in Q$  and  $\lambda\varphi + (1 - \lambda)\psi \in F$  for some  $\lambda$  strictly between 0 and 1. Weak\* closed faces of  $Q(A)$  are in one-to-one correspondence with closed projections in  $A^{**}$ . In fact, a projection  $p$  in  $A^{**}$  is *closed* if and only if the face

$$F(p) = \{\varphi \in Q(A) : \varphi(1 - p) = 0\}$$

of  $Q(A)$  supported by  $p$  is weak\* closed [16]. Closed projections  $p$  in  $A^{**}$  are also in one-to-one correspondence with norm closed left ideals  $L$  of  $A$  such that

$$L = A^{**}(1 - p) \cap A.$$

Moreover, we have isometrical isomorphisms  $a + L \mapsto ap$  and  $x + L^{**} \mapsto xp$  under which

$$A/L \cong Ap \quad \text{and} \quad (A/L)^{**} \cong A^{**}/L^{**} \cong A^{**}p$$

as Banach spaces, respectively [11, 17, 1].

In the following,  $p$  is always a closed projection in  $A^{**}$ . We consider  $Ap$  as the left quotient  $A/L$  of the  $C^*$ -algebra  $A$  by the norm closed left ideal  $L = A^{**}(1 - p) \cap A$ . Consequently, its Banach double dual  $A^{**}p$  is the left quotient  $A^{**}/L^{**}$  of the  $w^*$ -algebra  $A^{**}$  by the weak\* closed left ideal  $L^{**} = A^{**}(1 - p)$ .

**Proposition 1.4.** *Let  $x \in A^{**}$ .  $x$  has a continuous atomic part modulo  $L^{**}$  if and only if  $zxp = zap$  for some  $a$  in  $A$ .*

PROOF. It suffices to observe the following equivalences:  $zx + L^{**} = za + L^{**}$  if and only if  $z(x - a) \in L^{**}$  if and only if  $z(x - a)p = 0$ .  $\square$

We also have a function representation of left quotients.

**Theorem 1.5** ([20]). *Let  $xp \in A^{**}p$ .  $xp \in Ap$  if and only if the bounded and affine functionals  $\varphi \mapsto \varphi(x^*x)$  and  $\varphi \mapsto \varphi(a^*x)$ ,  $\forall a \in A$ , are continuous on  $F(p)$ .*

Note that the extreme boundary of  $F(p)$  is  $(P(A) \cup \{0\}) \cap F(p)$ , i.e. the set of all pure states of  $A$  supported by  $p$  together with the zero functional. Motivated by Theorem 1.1, we shall attack the following

**Problem 1.6.** *Let  $x \in A^{**}$ . Suppose that  $x^*x$  and  $a^*x$ ,  $\forall a \in A$ , are uniformly continuous on  $(P(A) \cup \{0\}) \cap F(p)$ . Can we infer that  $x$  has a continuous atomic part modulo  $L^{**}$ , i.e.  $zap \in zAp$ ?*

Problem 1.6 does not always have an affirmative answer (see counter examples in Section 4). Our main result, proved in Section 2, states

**Theorem 1.7.** *Let  $x \in A^{**}$ .  $x$  has a continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in zAp$ ) if and only if  $zpx^*xp \in zpAp$  and  $zpa^*xp \in zpAp$ ,  $\forall a \in A$ .*

An answer to Problem 1.6 should thus assert conditions on either  $x$  or  $p$  under which uniform continuities of  $x^*x$  and  $a^*x$ ,  $\forall a \in A$ , on the extreme boundary of  $F(p)$  will ensure  $zpx^*xp \in zpAp$  and  $zpa^*xp \in zpAp$ ,  $\forall a \in A$ . We shall see (via Theorem 2.4) that this amounts to looking for conditions under which uniformly continuous functionals of the extreme boundary of  $F(p)$  can be lifted to continuous affine functionals of the whole of  $F(p)$ . In Section 3, we shall present some sufficient conditions to ensure the existence of such liftings, and thus Problem 1.6 can be solved affirmatively. One of them assumes the universal measurability of  $x^*x$  and  $a^*x$  on  $F(p)$ . The others assume that  $p$  has either MSQC (this holds if, in particular,  $p$  is central) or  $p$  is semi-atomic. Counter examples in Section 4 verify that our conclusions are sharp.

## 2. MAIN RESULTS

Recall the notion  $(T, \{H_t\}, \Gamma)$  of a continuous field of Hilbert spaces [12, 10]. The *base space*  $T$  is a Hausdorff space, the *fiber*  $H_t$  is a Hilbert space (possibly different) for each  $t$  in  $T$  and the *continuous structure*  $\Gamma$  is a family of vector sections  $a = (a_t)_{t \in T}$  in the product space  $\prod_{t \in T} H_t$  satisfying the following two conditions.

1. The norm  $t \mapsto \|a_t\|_{H_t}$  is continuous on  $T$  for each  $a$  in  $\Gamma$ .
2. The set  $\{a_t : a \in \Gamma\}$  is norm dense in  $H_t$  for each  $t$  in  $T$ .

A vector section  $x = (x_t)_{t \in T}$  in  $\prod_{t \in T} H_t$  is said to be *bounded* if the functional  $t \mapsto \langle x_t, x_t \rangle_{H_t} = \|x_t\|_{H_t}^2$  is bounded on  $T$ . A bounded vector section  $x$  is said to be *weakly continuous* if the functionals  $t \mapsto \langle x_t, a_t \rangle_{H_t}$  is continuous on  $T$  for every  $a = (a_t)_{t \in T}$  in  $\Gamma$ . A weakly continuous vector section  $x$  is said to be *continuous* if, in addition, the functional  $t \mapsto \langle x_t, x_t \rangle_{H_t} = \|x_t\|_{H_t}^2$  is continuous. The subspaces of bounded, weakly continuous, and continuous vector sections in the product space  $\prod_{t \in T} H_t$  equipped with the sup norm  $\|x\|_\infty = \sup_{t \in T} \|x_t\|_{H_t}$  are Banach spaces, respectively.

Although the following elementary result should have been known, we provide a proof here since we cannot find any in the literature.

**Lemma 2.1.** *Let  $(T, \{H_t\}_t, \Gamma)$  be a continuous field of Hilbert spaces. Let  $(y_t)_{t \in T}$  be a weakly continuous vector section. Then the map  $t \mapsto \|y_t\|_{H_t}$  is lower semicontinuous on  $T$ .*

PROOF. For each  $t_0$  in  $T$ , since  $\{u_{t_0} \in H_{t_0} : u \in \Gamma\}$  is dense in  $H_{t_0}$ , we have

$$\|y_{t_0}\|_{H_{t_0}} = \sup\{|\langle y_{t_0}, u_{t_0} \rangle_{H_{t_0}}| : \|u_{t_0}\|_{H_{t_0}} < 1, u \in \Gamma\}.$$

For  $\varepsilon > 0$ , choose an  $a$  in  $\Gamma$  so that  $\|a_{t_0}\|_{H_{t_0}} < 1$  and

$$\|y_{t_0}\|_{H_{t_0}} - \langle y_{t_0}, a_{t_0} \rangle_{H_{t_0}} < \varepsilon/2.$$

Since  $t \mapsto \|a_t\|_{H_t}$  and  $t \mapsto \langle y_t, a_t \rangle_{H_t}$  are continuous on  $T$ , for those  $t$  close enough to  $t_0$  we have  $\|a_t\|_{H_t} < 1$  and

$$|\langle y_t, a_t \rangle_{H_t} - \langle y_{t_0}, a_{t_0} \rangle_{H_{t_0}}| < \varepsilon/2.$$

Now, if  $t_\alpha \rightarrow t_0$  in  $T$ , for  $\alpha$  large enough,

$$\begin{aligned} \|y_{t_\alpha}\|_{H_{t_\alpha}} &= \sup\{|\langle y_{t_\alpha}, u_{t_\alpha} \rangle_{H_{t_\alpha}}| : \|u_{t_\alpha}\|_{H_{t_\alpha}} < 1, u \in \Gamma\} \\ &\geq |\langle y_{t_\alpha}, a_{t_\alpha} \rangle_{H_{t_\alpha}}| \\ &> |\langle y_{t_0}, a_{t_0} \rangle_{H_{t_0}}| - \varepsilon/2 \\ &> \|y_{t_0}\|_{H_{t_0}} - \varepsilon. \end{aligned}$$

Consequently,

$$\|y_{t_0}\|_{H_{t_0}} \leq \liminf \|y_{t_\alpha}\|_{H_{t_\alpha}} + \varepsilon.$$

Since  $\varepsilon$  is arbitrary,

$$\|y_{t_0}\|_{H_{t_0}} \leq \liminf \|y_{t_\alpha}\|_{H_{t_\alpha}},$$

and thus  $t \mapsto \|y_t\|_{H_t}$  is lower semicontinuous on  $T$ .  $\square$

Let  $A$  be a  $C^*$ -algebra and  $L$  a norm closed left ideal of  $A$ . Let  $p$  be the closed projection in  $A^{**}$  related to  $L$  such that  $L = A^{**}(1-p) \cap A$ . Moreover,  $A/L \cong Ap$  and  $(A/L)^{**} \cong A^{**}/L^{**} \cong A^{**}p$  as Banach spaces. We are going to construct a continuous field of Hilbert spaces associated to the left quotient  $Ap$ . The base space is  $F(p)$ , the weak\* closed face of the quasi-state space  $Q(A)$  of  $A$  supported by  $p$ . Note that

$$\begin{aligned} F(p) &= \{\varphi \in Q(A) : \varphi(1-p) = 0\} \\ &= \{\varphi \in Q(A) : \varphi(x) = \varphi(xp) = \varphi(px) = \varphi(pxp), \forall x \in A^{**}\}. \end{aligned}$$

Moreover,  $F(p)$  is itself a weak\* compact, Hausdorff and convex set.

For each  $\varphi$  in  $F(p)$ , the GNS construction yields a cyclic representation  $(\pi_\varphi, H_\varphi, \omega_\varphi)$  of  $A$ . In particular,  $\varphi(x) = \langle \pi_\varphi(x)\omega_\varphi, \omega_\varphi \rangle_\varphi, \forall x \in A^{**}$ , where  $\langle \cdot, \cdot \rangle_\varphi$  is the inner product of the Hilbert space  $H_\varphi$ . Set  $H_\varphi$  to be the zero dimensional Hilbert space when  $\varphi = 0$ . Note also that for each  $\varphi$  in  $F(p)$ ,  $\pi_\varphi(p)\omega_\varphi = \omega_\varphi$  since  $\langle \pi_\varphi(p)\omega_\varphi, \omega_\varphi \rangle_\varphi = \varphi(p) = \varphi(1) = \|\omega_\varphi\|^2$ .

**Notation.** Write  $x\omega_\varphi$  for  $\pi_\varphi(x)\omega_\varphi$  in  $H_\varphi, \forall x \in A^{**}, \forall \varphi \in F(p)$ .

In this way, there is an embedding  $A^{**}p \hookrightarrow \prod_{\varphi \in F(p)} H_\varphi$  defined by associating each  $xp$  in  $A^{**}p$  to the vector section  $(x\omega_\varphi)_{\varphi \in F(p)}$  in  $\prod_{\varphi \in F(p)} H_\varphi$ . A continuous structure of  $\prod_{\varphi \in F(p)} H_\varphi$  can be defined by the image of  $Ap$  under this embedding. In fact, the functional  $\varphi \mapsto \|a\omega_\varphi\|^2 = \langle a\omega_\varphi, a\omega_\varphi \rangle_\varphi = \varphi(a^*a)$  is continuous on  $F(p)$  for every  $ap$  in  $Ap$ . Moreover,  $\pi_\varphi(A)\omega_\varphi = \{a\omega_\varphi : a \in A\}$  is norm dense in  $H_\varphi$ .

In the continuous field  $(F(p), \{H_\varphi\}, Ap)$  of Hilbert spaces, a bounded vector section  $(x_\varphi)_{\varphi \in F(p)}$  in the product space  $\prod_{\varphi \in F(p)} H_\varphi$  is weakly continuous if  $\varphi \mapsto \langle x_\varphi, a\omega_\varphi \rangle_\varphi$  is continuous on  $F(p)$  for all  $a$  in  $A$ . A weakly continuous vector section  $(x_\varphi)_{\varphi \in F(p)}$  is continuous if, in addition,  $\varphi \mapsto \langle x_\varphi, x_\varphi \rangle_\varphi$  is continuous on  $F(p)$ . Since  $F(p)$  is a convex set, we have an additional affine structure.

**Definition 2.2.** Let  $(x_\varphi)_{\varphi \in F(p)}$  be a vector section in  $\prod_{\varphi \in F(p)} H_\varphi$ .  $(x_\varphi)_{\varphi \in F(p)}$  is said to be *affine* if the functionals  $\varphi \mapsto \langle x_\varphi, a\omega_\varphi \rangle_\varphi$ ,  $\forall a \in A$ , are affine on  $F(p)$ . In other words,

$$\langle x_\varphi, a\omega_\varphi \rangle_\varphi = \lambda \langle x_\psi, a\omega_\psi \rangle_\psi + (1 - \lambda) \langle x_\rho, a\omega_\rho \rangle_\rho.$$

whenever  $\varphi = \lambda\psi + (1 - \lambda)\rho$  in  $F(p)$ ,  $0 \leq \lambda \leq 1$  and  $a \in A$ .

In [20], we showed that every bounded and affine vector section  $(x_\varphi)_{\varphi \in F(p)}$  arises from an  $xp$  in  $A^{**}p$ , i.e.  $x_\varphi = x\omega_\varphi$ ,  $\forall \varphi \in F(p)$ . More precisely, we have

**Theorem 2.3** ([20]). *Let  $A$  be a C\*-algebra and  $p$  a closed projection in  $A^{**}$ .*

1.  $A^{**}p$  is isometrically linear isomorphic to the Banach space of all bounded and affine vector sections in  $\prod_{\varphi \in F(p)} H_\varphi$  equipped with the norm  $\sup_{\varphi \in F(p)} \|x\omega_\varphi\|_{H_\varphi} = \sup_{\varphi \in F(p)} \varphi(x^*x)^{1/2}$ .
2.  $Ap$  is isometrically linear isomorphic to the Banach space of all continuous and affine vector sections in  $\prod_{\varphi \in F(p)} H_\varphi$ .

Theorem 1.5 is a corollary of Theorem 2.3. In fact,  $xp$  in  $A^{**}p$  defines a continuous vector section in  $\prod_{\varphi \in F(p)} H_\varphi$  if and only if the affine functionals  $\varphi \mapsto \langle x\omega_\varphi, x\omega_\varphi \rangle_\varphi = \varphi(x^*x)$  and  $\varphi \mapsto \langle x\omega_\varphi, a\omega_\varphi \rangle_\varphi = \varphi(a^*x)$ ,  $\forall a \in A$ , are continuous on  $F(p)$ . The meaning of this translation can be more precise with the help of the following result of Brown, where  $A_{sa}$  (resp.  $A_{sa}^{**}$ ) is the self-adjoint part of  $A$  (resp.  $A^{**}$ ).

**Theorem 2.4** ([6, 3.9]).  *$pA_{sa}p$  (resp.  $pA_{sa}^{**}p$ ) is isometrically linear and order isomorphic to the Banach space of all continuous (resp. bounded) affine functionals of  $F(p)$  which vanish at zero.*

Let  $\mathcal{W}_p$  denote the Banach subspace of  $A^{**}p$  of all weakly continuous affine vector sections in  $\prod_{\varphi \in F(p)} H_\varphi$ . When  $p = 1$ ,  $\mathcal{W}_p$  coincides with the set  $RM(A)$  of right multipliers of  $A$ . However,  $\mathcal{W}_p \neq RM(A)p$  in general [20].

In summary, Theorems 2.3 and 2.4 imply

**Corollary 2.5.** *Let  $xp \in A^{**}$ .*

1. *The following are all equivalent.*

- (a)  $xp \in \mathcal{W}_p$ .
  - (b) The affine functionals  $\varphi \mapsto \varphi(a^*x) = \varphi(pa^*xp)$ ,  $\forall a \in A$ , are continuous on  $F(p)$ .
  - (c)  $pa^*xp \in pAp$ ,  $\forall a \in A$ .
2. The following are all equivalent.
- (a)  $xp \in Ap$ .
  - (b) The affine functionals  $\varphi \mapsto \varphi(x^*x) = \varphi(px^*xp)$  and  $\varphi \mapsto \varphi(a^*x) = \varphi(pa^*xp)$ ,  $\forall a \in A$ , are continuous on  $F(p)$ .
  - (c)  $px^*xp \in pAp$  and  $pa^*xp \in pAp$ ,  $\forall a \in A$ .

Recall that  $xp$  in  $A^{**}p$  is said to have a continuous (resp. a weakly continuous) atomic part modulo  $L^{**}$  if  $zxp = zyp$  for some  $yp$  in  $Ap$  (resp. in  $\mathcal{W}_p$ ).

**Lemma 2.6.** *Let  $x, y \in A^{**}$ .  $zxp = zyp$  if and only if  $x\omega_\varphi = y\omega_\varphi$  in  $H_\varphi$  for every pure state  $\varphi$  in  $F(p)$ .*

PROOF. Let  $zxp = zyp$ . Since  $\pi_\varphi(z) = 1$  and  $\pi_\varphi(p)\omega_\varphi = \omega_\varphi$ ,  $x\omega_\varphi = \pi_\varphi(x)\omega_\varphi = \pi_\varphi(zxp)\omega_\varphi = \pi_\varphi(zyp)\omega_\varphi = \pi_\varphi(y)\omega_\varphi = y\omega_\varphi$  for every pure state  $\varphi$  in  $F(p)$ . For the converse, we note that for every  $\psi$  in  $Q(A)$ , the atomic  $\psi(zp \cdot p)$  can be written as a countable sum of pure positive linear functionals in  $F(p)$  [13]. Now,  $x\omega_\varphi = y\omega_\varphi$  implies  $\varphi((x-y)^*(x-y)) = \langle (x-y)\omega_\varphi, (x-y)\omega_\varphi \rangle_\varphi = 0$ . If this holds for all pure states  $\varphi$  in  $F(p)$  then  $\psi(zp(x-y)^*(x-y)p) = 0$  for all  $\psi$  in  $Q(A)$ . Consequently,  $zp(x-y)^*(x-y)p = 0$  and thus  $zxp = zyp$ .  $\square$

**Theorem 2.7.** *Let  $x \in A^{**}$ .  $x$  has a weakly continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in z\mathcal{W}_p$ ) if and only if  $zpa^*xp \in zpAp$  for all  $a$  in  $A$ .*

PROOF. For the necessity, we assume that  $zxp = zyp$  for some  $yp$  in  $\mathcal{W}_p$ . Then  $pa^*yp \in pAp$  by Corollary 2.5, and thus  $zpa^*xp = pa^*(zxp) = pa^*(zyp) = z(pa^*yp) \in zpAp$ ,  $\forall a \in A$ .

For the sufficiency, let  $X = F(p) \cap P(A)$ , the set of all pure states of  $A$  supported by  $p$ . We want to find a  $yp$  in  $\mathcal{W}_p$  such that  $x\omega_\varphi = y\omega_\varphi$  for all  $\varphi$  in  $X$ . In this case, we shall have  $zxp = zyp \in z\mathcal{W}_p$  by Lemma 2.6.

Since pure states are also supported by the central projection  $z$ , we have

$$\varphi(x) = \varphi(pxp) = \varphi(zpxp), \quad \forall x \in A^{**}, \forall \varphi \in X.$$

For each  $a$  in  $A$ , by the hypothesis,  $zpa^*xp = zpv_a p$  for some  $v_a$  in  $A$ . And thus

$$(1) \quad \langle x\omega_\varphi, a\omega_\varphi \rangle_\varphi = \varphi(a^*x) = \varphi(zpa^*xp) = \varphi(zpv_a p) = \varphi(v_a), \quad \forall \varphi \in X.$$

Let  $\overline{X}$  be the weak\* closure of  $X$  in  $F(p)$ . If  $\psi_\alpha \in X$  and  $\psi = \lim \psi_\alpha \in \overline{X}$  then we have, by (1),

$$(2) \quad |\psi(v_a)| = \lim |\psi_\alpha(v_a)| = \lim |\langle x\omega_{\psi_\alpha}, a\omega_{\psi_\alpha} \rangle_{\psi_\alpha}| \\ \leq \|x\| \lim \|a\omega_{\psi_\alpha}\|_{\psi_\alpha} = \|x\| \|a\omega_\psi\|_\psi = \|x\| \|\psi(a^*a)\|^{1/2}.$$

Note that  $X \cup \{0\}$  is the extreme boundary of the compact convex set  $F(p)$ . Thus each continuous affine functionals of  $F(p)$  assumes its maximum modules at a point in  $X$ . From

Theorem 2.4 we know that there is an order-preserving linear isometry from  $pA_{sa}p$  into the Banach space  $C_{\mathbb{R}}(\overline{X})$  of continuous real-valued functions defined on  $\overline{X}$ . Hence, as a positive linear functional of  $pA_{sa}p$ , each  $\varphi$  in  $F(p)$  has a (non-unique) Hahn-Banach positive extension  $m_{\varphi}$  in the space  $M(\overline{X})$  of regular finite Borel measures on the compact Hausdorff space  $\overline{X}$ . Thus we can write

$$(3) \quad \varphi(a) = \varphi(pap) = \int_{\overline{X}} \psi(pap) dm_{\varphi}(\psi) = \int_{\overline{X}} \psi(a) dm_{\varphi}(\psi), \quad \forall a \in A_{sa}, \forall \varphi \in F(p).$$

Motivating by (1), we define a vector section  $(y_{\varphi})_{\varphi \in F(p)}$  in  $\prod_{\varphi \in F(p)} H_{\varphi}$  by the conditions

$$\langle y_{\varphi}, a\omega_{\varphi} \rangle_{\varphi} = \varphi(v_a) = \int_{\overline{X}} \psi(v_a) dm_{\varphi}(\psi), \quad \forall a \in A, \forall \varphi \in F(p).$$

Observe that

$$\begin{aligned} |\langle y_{\varphi}, a\omega_{\varphi} \rangle_{\varphi}| &\leq \int_{\overline{X}} |\psi(v_a)| dm_{\varphi}(\psi) \\ &\leq \int_{\overline{X}} \|x\| \psi(a^*a)^{1/2} dm_{\varphi}(\psi) \quad (\text{by (2)}) \\ &\leq \|x\| \left[ \int_{\overline{X}} \psi(a^*a) dm_{\varphi}(\psi) \right]^{1/2} \quad (\text{since } m_{\varphi}(\overline{X}) = \|\varphi\| \leq 1) \\ &= \|x\| \varphi(a^*a)^{1/2} \quad (\text{by (3)}) \\ &= \|x\| \|a\omega_{\varphi}\|_{\varphi}. \end{aligned}$$

Therefore, the definition of  $y_{\varphi}$  (as a bounded linear functional of the Hilbert space  $H_{\varphi}$ ) makes sense and  $\|(y_{\varphi})_{\varphi \in F(p)}\|_{\infty} = \sup_{\varphi \in F(p)} \|y_{\varphi}\|_{H_{\varphi}} \leq \|x\|$ . Clearly, the definition of  $y_{\varphi}$  is independent of the choice of  $m_{\varphi}$  and  $v_a$ . Since  $\varphi \mapsto \langle y_{\varphi}, a\omega_{\varphi} \rangle_{\varphi} = \varphi(v_a)$  is a continuous affine functional of  $F(p)$  for each  $a$  in  $A$ ,  $(y_{\varphi})_{\varphi \in F(p)}$  is a bounded and weakly continuous affine vector section in  $\prod_{\varphi \in F(p)} H_{\varphi}$ . By Theorem 2.3, there is a  $yp$  in  $\mathcal{W}_p$  such that  $y\omega_{\varphi} = y_{\varphi}$  for each  $\varphi$  in  $F(p)$ . Finally, for each pure state  $\varphi$  in  $F(p)$  and  $a$  in  $A$  we have  $\langle y\omega_{\varphi}, a\omega_{\varphi} \rangle_{\varphi} = \varphi(v_a) = \langle x\omega_{\varphi}, a\omega_{\varphi} \rangle_{\varphi}$  by (1). Since  $\{a\omega_{\varphi} : a \in A\}$  is norm dense in (indeed, equal to)  $H_{\varphi}$ ,  $y\omega_{\varphi} = x\omega_{\varphi}$  for every pure state  $\varphi$  in  $F(p)$ . Thus,  $zxp = zyp$ .  $\square$

Beside Theorem 1.3, the following result supplements Theorem 1.1 in another interesting way. Note that  $\mathcal{W}_p = RM(A)$  when  $p = 1$  (Corollary 2.5).

**Corollary 2.8.** *Let  $x \in A^{**}$ .  $zx = zy$  for some right multiplier  $y$  of  $A$  in  $A^{**}$  if and only if  $a^*y, \forall a \in A$ , are uniformly continuous on  $P(A) \cup \{0\}$ .*

Now we are ready to present the proof of our main result, Theorem 1.7, which says that for each  $x$  in  $A^{**}$ ,  $x$  has a continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in zAp$ ) if and only if  $zpx^*xp \in zpAp$  and  $zpa^*xp \in zpAp, \forall a \in A$ .

**PROOF OF THEOREM 1.7.** Only the sufficiency demands a proof. By Theorem 2.7,  $x$  has a weakly continuous atomic part modulo  $L^{**}$ , i.e.  $zxp = zyp$  for some  $yp$  in  $\mathcal{W}_p$ . By hypothesis, we also have  $zpx^*xp = zpvv$  for some  $v$  in  $A$ . Because  $pA_+p = (pAp)_+$ , we can assume  $v$  to be positive. By Lemma 2.1,  $\varphi \mapsto \varphi(y^*y - v) = \|y\omega_{\varphi}\|_{H_{\varphi}}^2 - \varphi(v)$  is a lower semicontinuous

real-valued affine functional of the compact convex set  $F(p)$ , which vanishes at all pure states in  $F(p)$ . Now the desired assertion follows from a result of Pedersen [15, 3.8]. However, for the convenience of the readers we present below a somewhat elementary argument. We note that the scalar function  $\varphi \mapsto \varphi(y^*y - v)$  attains its minimum value at an extreme point  $\psi$  of  $F(p)$ , *i.e.* a pure state  $\psi$  in  $F(p)$ . But  $\psi(y^*y - v) = \psi(zp(y^*y - v)p) = 0$  for all pure states  $\psi$  in  $F(p)$ . Hence,  $py^*yp \geq pvp$ . On the other hand, it follows from the Krein-Milman Theorem that every  $\varphi$  in the compact convex set  $F(p)$  is a weak\* limit of convex combinations  $\varphi_\alpha$  of pure states (and 0) in  $F(p)$ . Therefore, we always have  $\varphi(y^*y) \leq \liminf \varphi_\alpha(y^*y) = \liminf \varphi_\alpha(v) = \varphi(v)$  by Lemma 2.1. Thus,  $py^*yp \leq pvp$ . Consequently,  $py^*yp = pvp \in pAp$ . Hence,  $yp \in Ap$  by Corollary 2.5. Finally,  $zxp = zyp \in zAp$ .  $\square$

In Section 3, we shall investigate conditions under which our main result, Theorem 1.7, can be translated into an answer of Problem 1.6.

### 3. SOME APPROACHES TO THE PROBLEM

**3.1. Universally measurable elements .** Let  $A$  be a  $C^*$ -algebra and  $p$  a closed projection in  $A^{**}$ . Recall that  $A_{sa}^m$  consists of all limits in  $A_{sa}^{**}$  of monotone increasing nets in  $A_{sa}$  and  $(A_{sa})_m = -A_{sa}^m$ . While  $A_{sa}$  consists of continuous affine real-valued functionals of  $Q(A)$  (the Kadison function representation), the norm closure  $(A_{sa}^m)^-$  of  $A_{sa}^m$  consists of *lower semicontinuous elements* and the norm closure  $\overline{(A_{sa})_m}$  of  $(A_{sa})_m$  consists of *upper semicontinuous elements* in  $A^{**}$ . Accordingly, an element  $x$  of  $A_{sa}^{**}$  is said to be *universally measurable* if for each  $\varphi$  in  $Q(A)$  and  $\varepsilon > 0$  there exist a lower semicontinuous element  $l$  and an upper semicontinuous element  $u$  in  $A^{**}$  such that  $u \leq x \leq l$  and  $\varphi(l - u) < \varepsilon$  [15].

We note that  $pA_{sa}p$  consists of continuous affine real-valued functionals of the weak\* closed face  $F(p)$  of  $Q(A)$  supported by  $p$  (Theorem 2.4). Analogously,  $pxp$  in  $pA_{sa}^{**}p$  is said to be *universally measurable on  $F(p)$*  if for each  $\varphi$  in  $F(p)$  and  $\varepsilon > 0$ ,  $l$  in  $(A_{sa}^m)^-$  and  $u$  in  $\overline{(A_{sa})_m}$  exist such that  $pup \leq pxp \leq plp$  and  $\varphi(l - u) < \varepsilon$ .  $pxp$  in  $pA^{**}p$  is said to be *universally measurable on  $F(p)$*  if both the real and imaginary parts of  $pxp$  are.

**Lemma 3.1.** *Let  $x \in A_{sa}^{**}$  and  $\overline{X}$  the weak\* closure of  $X = F(p) \cap P(A)$  in  $F(p)$ . If  $pxp$  is universally measurable on  $F(p)$  and continuous on  $\overline{X}$  then  $pxp \in pAp$ .*

**PROOF.** First, we note that the continuity of  $pxp$  on  $\overline{X}$  means that whenever  $\varphi_\lambda \rightarrow \varphi$  weak\* in  $\overline{X}$  we have  $\varphi_\lambda(x) \rightarrow \varphi(x)$ . In view of Theorem 2.4, we need to verify these convergences for nets arising from the whole of  $F(p)$ .

Suppose there were  $\varphi_\lambda, \varphi$  in  $F(p)$  such that  $\varphi_\lambda \rightarrow \varphi$  weak\* but  $\varphi_\lambda(x)$  did not converge to  $\varphi(x)$ . Without loss of generality, we can assume there exists a  $\delta > 0$  such that  $|\varphi(x) - \varphi_\lambda(x)| > \delta$  for all  $\lambda$ . As in the proof of Theorem 2.7, we can embed  $pA_{sa}p$  as a closed subspace into the Banach space  $C_{\mathbb{R}}(\overline{X})$  of continuous real-valued functions defined on  $\overline{X}$ . Let  $m_\lambda$  be any positive extension of  $\varphi_\lambda$  from  $pA_{sa}p$  to  $C_{\mathbb{R}}(\overline{X})$  and  $\|m_\lambda\| = \|\varphi_\lambda\| \leq 1$ . Hence,  $(m_\lambda)_\lambda$  is a bounded net in  $M(\overline{X})$ , the Banach dual space of  $C_{\mathbb{R}}(\overline{X})$ , consisting of regular finite Borel measures on the

compact Hausdorff space  $\overline{X}$ . Then, by passing to a subnet if necessary, we have  $m_\lambda \rightarrow m$  in the weak\* topology of  $M(\overline{X})$ . Clearly,  $m \geq 0$  and  $m|_{pA_{sa}p} = \varphi$ . On the other hand,  $pxp$  is assumed to be universally measurable on  $F(p)$ . So for each  $\phi$  in  $F(p)$  and  $\varepsilon > 0$  there exist a  $u$  in  $(A_{sa})_m^-$  and an  $l$  in  $(A_{sa}^m)^-$  such that

$$pup \leq pxp \leq plp \quad \text{and} \quad \phi(l - u) < \varepsilon.$$

It follows from the semicontinuity and the affine property of  $u$  and  $l$  (cf. [5, p. 19], or by a direct argument) that  $u$  and  $l$  satisfy the barycenter formula of  $\phi$  in  $F(p)$ , *i.e.*

$$\phi(u) = \int_{\overline{X}} \psi(u) dm_\phi(\psi) \quad \text{and} \quad \phi(l) = \int_{\overline{X}} \psi(l) dm_\phi(\psi),$$

where  $m_\phi$  in  $M(\overline{X})$  is a positive Hahn-Banach extension of  $\phi$  to  $C_{\mathbb{R}}(\overline{X})$ . Since  $pup \leq pxp \leq plp$ , we have

$$\phi(u) = \int_{\overline{X}} \psi(u) dm_\phi(\psi) \leq \int_{\overline{X}} \psi(x) dm_\phi(\psi) \leq \int_{\overline{X}} \psi(l) dm_\phi(\psi) = \phi(l), \quad \forall \phi \in F(p).$$

As  $\phi(u) \leq \phi(x) \leq \phi(l)$ , we have

$$\left| \phi(x) - \int_{\overline{X}} \psi(x) dm_\phi(\psi) \right| \leq \phi(l - u) < \varepsilon, \quad \forall \phi \in F(p).$$

Because  $\varepsilon$  is arbitrary,  $x$  satisfies the barycenter formula of  $\phi$  in  $F(p)$  as well, *i.e.*

$$\phi(x) = \int_{\overline{X}} \psi(x) dm_\phi(\psi), \quad \forall \phi \in F(p).$$

Note again that  $\psi(x) = \psi(pxp)$  for each  $\psi$  in  $F(p)$ . Now  $pxp \in C_{\mathbb{R}}(\overline{X})$  implies

$$\int_{\overline{X}} \psi(x) dm_\lambda(\psi) = \int_{\overline{X}} \psi(pxp) dm_\lambda(\psi) \longrightarrow \int_{\overline{X}} \psi(pxp) dm(\psi) = \int_{\overline{X}} \psi(x) dm(\psi).$$

Consequently,  $\varphi_\lambda(x) \longrightarrow \varphi(x)$ , a contradiction.  $\square$

In the proof above, what we actually need on  $x$  is that  $pxp$  satisfies barycenter formulas of elements of  $F(p)$  and are continuous on  $\overline{X} = \overline{P(A) \cap F(p)}$ . Indeed, we have proved

**Proposition 3.2.** *For every  $pxp$  in  $pA^{**}p$  satisfying barycenter formulas of elements of  $F(p)$ ,  $pxp \in pAp$  if and only if  $pxp$  is continuous on  $\overline{X}$ .*

**Theorem 3.3.** *Let  $x \in A^{**}$  and  $X = F(p) \cap P(A)$  with weak\* closure  $\overline{X}$  in  $F(p)$ .*

1.  $xp \in \mathcal{W}_p$  if and only if  $pa^*xp, \forall a \in A$ , are universally measurable on  $F(p)$  and continuous on  $\overline{X}$ .
2.  $xp \in Ap$  if and only if  $px^*xp$  and  $pa^*xp, \forall a \in A$ , are universally measurable on  $F(p)$  and continuous on  $\overline{X}$ .

PROOF. We note that the real and imaginary parts of  $pa^*xp$  both satisfy the assumptions of Lemma 3.1 for any  $a$  in  $A$ . Hence, the assertions follow from Lemma 3.1 and Corollary 2.5.  $\square$

**3.2.  $p$  has MSQC .** Let  $A$  be a  $C^*$ -algebra. Recall that a projection  $p$  in  $A^{**}$  is closed if the face  $F(p) = \{\varphi \in Q(A) : \varphi(1-p) = 0\}$  of  $Q(A)$  supported by  $p$  is weak\* closed.  $p$  is said to be *compact* if  $F(p) \cap S(A)$  is closed [6], where  $S(A) = \{\varphi \in Q(A) : \|\varphi\| = 1\}$  is the state space of  $A$ . Let  $p$  be a closed projection in  $A^{**}$ .  $h$  in  $pA_{sa}^{**}p$  is said to be *q-continuous* on  $p$  [2] if the spectral projection  $E_F(h)$  (computed in  $pA^{**}p$ ) is closed for every closed subset  $F$  of  $\mathbb{R}$ .  $h$  is said to be *strongly q-continuous* on  $p$  [6] if, in addition,  $E_F(h)$  is compact whenever  $F$  is closed and  $0 \notin F$ . It is known from [6, 3.43] that  $h$  is strongly q-continuous on  $p$  if and only if  $h = pa = ap$  for some  $a$  in  $A_{sa}$ . In general,  $h$  in  $pA^{**}p$  is said to be *strongly q-continuous* if both  $\operatorname{Re}h$  and  $\operatorname{Im}h$  are.

Denote by  $SQC(p)$  the  $C^*$ -algebra of all strongly q-continuous elements on  $p$ . We say that  $p$  has MSQC (“many strongly q-continuous elements”) if  $SQC(p)$  is  $\sigma$ -weakly dense in  $pA^{**}p$ . Brown [8] showed that  $p$  has MSQC if and only if  $pAp = SQC(p)$  if and only if  $pAp$  is an algebra. In particular, every central projection  $p$  (especially,  $p = 1$ ) has MSQC. We provide a partial answer to Problem 1.6 by the following

**Theorem 3.4.** *Let  $p$  have MSQC and  $xp \in A^{**}p$ . Let  $X_0 = (F(p) \cap P(A)) \cup \{0\}$  be the extreme boundary of  $F(p)$ .*

1.  *$x$  has a weakly continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in z\mathcal{W}_p$ ) if and only if  $pa^*xp$ ,  $\forall a \in A$ , are uniformly continuous on  $X_0$ .*
2.  *$x$  has a continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in zAp$ ) if and only if  $px^*xp$  and  $pa^*xp$ ,  $\forall a \in A$ , are uniformly continuous on  $X_0$ .*

PROOF. The necessities are obvious and we check the sufficiencies. Note that  $pAp$  is now a  $C^*$ -algebra with the pure state space  $P(pAp) = F(p) \cap P(A)$ . The maximal atomic projection of  $pAp$  is  $zp$ . By Theorem 1.1,  $zpa^*xp \in zpAp$  for all  $a$  in  $A$ . In case  $px^*xp$  is uniformly continuous on  $X_0$ ,  $zpx^*xp \in zpAp$  as well. The assertion follows from Theorems 2.7 and 1.7.  $\square$

**Corollary 3.5.** *Let  $p$  have MSQC and  $xp \in A^{**}p$ . If  $pa^*xp$  is continuous on  $\overline{X} = \overline{F(p) \cap P(A)}$  for all  $a$  in  $A$  then  $zxp \in z\mathcal{W}_p$ . If, in addition,  $px^*xp$  is continuous on  $\overline{X}$  then  $zxp \in zAp$ .*

PROOF. We simply note that either 0 belongs to  $\overline{X}$  or 0 is isolated from  $X = F(p) \cap P(A)$  in  $F(p)$ . Consequently, continuity on the compact set  $\overline{X}$  ensures uniform continuity on  $X_0 = (F(p) \cap P(A)) \cup \{0\}$ . Thus, Theorem 3.4 applies.  $\square$

**3.3.  $p$  is semiatomic .** Let  $A$  be a  $C^*$ -algebra and  $p$  a closed projection in  $A^{**}$ . Recall that  $A$  is said to be scattered if  $Q(A) \subseteq zQ(A)$  [13, 14] and  $p$  is said to be atomic if  $F(p) \subseteq zF(p)$  [8]. If  $A$  is scattered then every closed projection in  $A^{**}$  is atomic. Moreover,  $A$  is said to be semiscattered [3] if  $\overline{P(A)} \subseteq zQ(A)$ .

Analogously, we say that a closed projection  $p$  is *semiatomic* if the weak\* closure of  $F(p) \cap P(A)$  contains only atomic positive linear functionals of  $A$ , i.e.  $\overline{F(p) \cap P(A)} \subseteq zF(p)$ . It is easy to see that if  $A$  is semiscattered then every closed projection in  $A^{**}$  is semiatomic. We provide another partial answer to Problem 1.6 by the following

**Theorem 3.6.** *Let  $p$  be semiatomic and  $xp \in A^{**}p$ . Let  $\overline{X} = \overline{F(p) \cap P(A)}$ .*

1.  *$x$  has a weakly continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in z\mathcal{W}_p$ ) if and only if  $pa^*xp$ ,  $\forall a \in A$ , are continuous on  $\overline{X}$ .*
2.  *$x$  has a continuous atomic part modulo  $L^{**}$  (i.e.  $zxp \in zAp$ ) if and only if  $px^*xp$  and  $pa^*xp$ ,  $\forall a \in A$ , are continuous on  $\overline{X}$ .*

To prove Theorem 3.6, we need the following generalization of [7, Theorem 6] in which  $p = 1$ .

**Lemma 3.7.** *Let  $x$  in  $zpA^{**}p$  be uniformly continuous on  $X_0 = (F(p) \cap P(A)) \cup \{0\}$ . Then  $x$  is in the C\*-algebra  $B$  generated by  $zpAp$ . In particular,  $x = zy$  for some universally measurable element  $y$  of  $pA^{**}p$ .*

PROOF. Utilizing a similar argument in [7], we can prove that  $x \in B$ . Moreover, the last assertion can be deduced from [4, 2.1] and the fact that  $\widetilde{B}_0$  is a Jordan algebra [9].  $\square$

Lemma 3.7 merely says that if  $x \in A^{**}$  such that  $x$  is uniformly continuous on the extreme boundary  $(F(p) \cap P(A)) \cup \{0\}$  of  $F(p)$  then  $x$  has a universally measurable atomic part modulo  $L^{**}$ .

PROOF OF THEOREM 3.6. We prove the sufficiencies only. Let  $xp$  in  $A^{**}p$  satisfy the stated conditions. For each  $a$  in  $A$ ,  $zpa^*xp$  is thus uniformly continuous on  $X_0 = (P(A) \cap F(p)) \cup \{0\}$ . By Lemma 3.7, there is a universally measurable element  $y$  of  $pA^{**}p$  such that  $zpa^*xp = zy$ . Since  $p$  is assumed to be semiatomic, each  $\varphi$  in  $\overline{X} = \overline{P(A) \cap F(p)}$  is atomic and thus  $\varphi(a^*x) = \varphi(zpa^*xp) = \varphi(zy) = \varphi(y)$ . In particular, the universally measurable element  $y$  is continuous on  $\overline{X}$ . It follows from Lemma 3.1 that  $y \in pAp$ . As a consequence,  $zpa^*xp \in zpAp$  for each  $a$  in  $A$ . It should be clear that we can also deduce  $zpx^*xp \in zpAp$  in the same manner if  $px^*xp$  is assumed to be continuous on  $\overline{X}$ . Now, Theorems 2.7 and 1.7 apply.  $\square$

#### 4. SOME COUNTER EXAMPLES

In this final section, we shall provide some counter examples to Problem 1.6 to show that the conclusions in Section 3 are sharp. Recall that  $X_0 = (P(A) \cap F(p)) \cup \{0\}$  and  $\overline{X} = \overline{F(p) \cap P(A)}$ .

First of all, Example 4.2 below tells us that the continuity assumptions in Lemma 3.1 on  $\overline{X}$  cannot be replaced by the uniform continuity on  $X_0$ . Moreover, we shall see in Example 4.1 that the conditions on universal measurability is also necessary. One may notice that uniform continuity of an element  $pxp$  in  $pA^{**}p$  on  $X_0$  ensures that  $pxp$  is universally measurable on  $X_0$  (Lemma 3.7). Unfortunately, even in this case (i.e.  $pxp$  is continuous on  $\overline{X}$  and thus universally measurable on  $X_0$ ) we can have  $zpxp \notin zpAp$  as shown in Example 4.3. Consequently, without assuming the universal measurability of  $px^*xp$  or  $pa^*xp$  on  $F(p)$ , Theorem 3.3 fails to give us any new result about the atomic part of  $x$ .

We now turn our attention to the MSQC assumption on the closed projection  $p$ . If  $p$  does not have MSQC then the conclusion of Theorem 3.4 may not hold. In fact, without assuming  $p$  has MSQC we have counter-examples; in one of which  $px^*xp$  and  $pa^*xp$ ,  $\forall a \in A$ , are universally

measurable on  $F(p)$  and uniformly continuous on  $X_0$  (Example 4.2), and in the other one  $px^*xp$  and  $pa^*xp, \forall a \in A$ , are continuous on  $\overline{X}$  (Example 4.3). But  $zxp \notin z\mathcal{W}_p$  in both cases.

Finally, we shall see in Example 4.2 that the continuity assumption on  $\overline{X}$  in Theorem 3.6 cannot be replaced by the uniform continuity on  $X_0$  even when the C\*-algebra  $A$  is scattered. On the other hand, Example 4.3 tells us that the conclusion of Theorem 3.6 can be wrong if  $p$  is not semiatomic.

**Example 4.1.** This example tells us that measurability conditions in Theorem 3.3 are necessary.

Let  $A = C(Y)$ , the abelian C\*-algebra of continuous (complex-valued) functions defined on a compact Hausdorff space  $Y$ .  $A^* \cong \bigoplus_1 \{L^1(\mu) : \mu \in \mathcal{C}\} \oplus_1 \ell^1(Y)$ , where  $\mathcal{C}$  is a maximal family of mutually singular continuous measures on  $Y$  of total variation one. Accordingly,  $A^{**} = \bigoplus_\infty \{L^\infty(\mu) : \mu \in \mathcal{C}\} \oplus_\infty \ell^\infty[0, 1]$ . Let  $p = 1$  in  $A^{**}$ . Pure states of  $A$  are exactly evaluations at points in  $[0, 1]$ . The maximal atomic projection  $z = 0 \oplus 1$  in the above direct sum decomposition of  $A^{**}$ . The embedding of  $A$  into  $A^{**}$  is given by  $g \mapsto (\bigoplus_{\mu \in \mathcal{C}} g_\mu) \oplus g_a$ , where  $g_\mu = g_a, \mu$  almost everywhere in  $Y$ , for all  $\mu$  in  $\mathcal{C}$ , and  $f_a = f$  everywhere in  $Y$ . Write  $f = f_d + f_a \in A^{**}$ , where  $f_d$  is the diffuse part of  $f$  which comes from  $\bigoplus_\infty \{L^\infty(\mu) : \mu \in \mathcal{C}\}$  and  $f_a$  is the atomic part of  $f$  which comes from  $\ell^\infty(Y)$ . Set  $f_d = 0$  and  $f_a$  to coincide with any nonzero continuous function on  $Y$ .  $f$  is *not* universally measurable (and neither satisfies barycenter formulas of elements of  $Q(A)$ ). Although  $f$  is continuous on  $\overline{P(A)} \cong Y$ ,  $f$  does not belong to  $A$ .  $\square$

**Example 4.2.** This example tells us that  $p$  having MSQC is necessary in Theorem 3.4 and continuity on  $\overline{X}$  is necessary in Theorem 3.6.

Let  $A$  be the scattered C\*-algebra of sequences of  $2 \times 2$  matrices  $x = (x_n)_{n=1}^\infty$  such that

$$x_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix} \longrightarrow x_\infty = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \text{ entrywise,}$$

and equipped with the  $\ell^\infty$ -norm. Note that the maximal atomic projection  $z = 1$  in this case. Let

$$p_n = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, n = 1, 2, \dots, \quad \text{and} \quad p_\infty = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then  $p = (p_n)_{n=1}^\infty$  is a closed projection in  $A^{**}$ . An element  $xp = (x_n p_n)_{n=1}^\infty$  of  $A^{**}p$  with

$$x_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}, n = 1, 2, \dots, \quad \text{and} \quad x_\infty = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix},$$

belongs to  $Ap$  if and only if  $a_n + b_n \rightarrow a$  and  $c_n + d_n \rightarrow d$ . Moreover,  $\mathcal{W}_p = Ap$  in this case.

We claim that  $p$  does *not* have MSQC. In fact, suppose  $x = (x_n)_{n=1}^\infty$  in  $A$  is given by

$$x_n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}, n = 1, 2, \dots, \quad \text{and} \quad x_\infty = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$$

such that  $x_n \rightarrow x_\infty$ . Then  $(pxp)_n = \lambda_n p_n, n = 1, 2, \dots$ , and  $(pxp)_\infty = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$  where  $\lambda_n = \frac{a_n + b_n + c_n + d_n}{2} \rightarrow \frac{a+d}{2}$ . Consequently,  $(pxp)_n^2 = \lambda_n^2 p_n, n = 1, 2, \dots$ , and  $(pxp)_\infty^2 = \begin{pmatrix} a^2 & 0 \\ 0 & d^2 \end{pmatrix}$ . If

$(pxp)^2 \in pAp$  we must have  $\lambda_n^2 \rightarrow \frac{a^2+d^2}{2}$ . This occurs exactly when  $a = d$ . In particular,  $pAp$  is not an algebra and thus  $p$  does *not* have MSQC.

On the other hand, the set  $X = P(A) \cap F(p)$  of all pure states in  $F(p)$  consists exactly of  $\varphi_n$ ,  $\psi_1$  and  $\psi_2$  which are given by

$$\varphi_n(x) = \text{tr}(x_n p_n), \quad n = 1, 2, \dots,$$

and

$$\psi_1(x) = a, \quad \psi_2(x) = d,$$

where  $x = (x_n)_{n=1}^\infty \in A$  and  $x_\infty = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$ . Since  $\varphi_n \rightarrow \frac{1}{2}(\psi_1 + \psi_2) \neq 0$ ,  $X_0 = X \cup \{0\}$  is discrete. Consider  $yp = (y_n p_n)_{n=1}^\infty$  in  $A^{**}p$  given by

$$y_n = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad n = 1, 2, \dots, \quad \text{and} \quad y_\infty = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Now, the universally measurable elements  $pa^*yp$  and  $px^*xp$  are uniformly continuous on  $X_0$  for all  $a$  in  $A$  but  $yp \notin Ap = \mathcal{W}_p$ .  $\square$

**Example 4.3.** This example is based on the ones in [18] and [3]. It tells us that without assuming  $p$  has MSQC or  $p$  is semiatomic, continuity on  $\overline{X}$  cannot ensure  $zxp \in zAp$ .

Let  $A$  be a C\*-algebra given by the exact sequence

$$0 \longrightarrow \bigoplus_{c_0} M_n \longrightarrow A \longrightarrow C[0, 1] \longrightarrow 0,$$

where  $M_n$  is the C\*-algebra of  $n \times n$  matrices,  $n = 1, 2, \dots$ . More precisely, if

$$h_n = \begin{pmatrix} 1/n & & & 0 \\ & 2/n & & \\ & & \ddots & \\ 0 & & & n/n \end{pmatrix}_{n \times n}, \quad n = 1, 2, \dots,$$

then we can implement  $A$  as the family of bounded sequences  $a = (a_n)_{n=1}^\infty$  such that  $a_\infty = f \in C[0, 1]$ ,  $a_n \in M_n$ ,  $n = 1, 2, \dots$ , and  $\|a_n - f(h_n)\| \rightarrow 0$ . Moreover,  $A^{**} \cong (\bigoplus_{\ell^\infty} M_n) \oplus_{\ell^\infty} C[0, 1]^{**}$ .

Let  $p_n$  be the projection  $\frac{1}{n} \begin{pmatrix} 1 & 1 & \dots & 1 \\ 1 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix}_{n \times n}$  in  $M_n$ ,  $n = 1, 2, \dots$ , and  $p_\infty = 1$  in  $C[0, 1]^{**}$ .

It is not difficult to see that  $p = (p_n)_{n=1}^\infty$  is a closed projection in  $A^{**}$  but  $p$  does *not* have MSQC. Moreover,  $X = P(A) \cap F(p) = \{\varphi_n : n = 1, 2, \dots\} \cup \{\chi_t : t \in [0, 1]\}$ . Here  $\varphi_n(a) = \text{tr}(a_n p_n)$ ,  $n = 1, 2, \dots$ , and  $\chi_t(a) = f(t)$ ,  $t \in [0, 1]$ , for every  $a = (a_n)_{n=1}^\infty$  in  $A$  with  $a_\infty = f$  in  $C[0, 1]$ . Since

$$\text{tr}(f(h_n)p_n) = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) \longrightarrow \int_0^1 f(t) dt$$

for every  $f$  in  $C[0, 1]$ , we have

$$\varphi_n(a) \longrightarrow \int_0^1 f(t) dt$$

for every  $a = (a_n)_{n=1}^\infty$  in  $A$  with  $a_\infty = f$  in  $C[0, 1]$ . Let  $\varphi_\infty = \lim \varphi_n$ . Note that for  $x = (x_n)_{n=1}^\infty$  in  $A^{**}$ ,  $\varphi_\infty(x) = \int_0^1 g_m(t) dt$  where  $x_\infty \in C[0, 1]^{**}$  and  $g_m$  is the component of  $x_\infty$  in  $L^\infty([0, 1], m)$  for the Lebesgue measure  $m$  on  $[0, 1]$  (cf. Example 4.1). It is then obvious that  $\overline{X} = \overline{P(A) \cap F(p)} = X \cup \{\varphi_\infty\}$ . Since  $\varphi_\infty$  is diffuse,  $p$  is *not* semiatomic.

Consider the element  $x = (x_n)_{n=1}^\infty$  of  $A^{**}$  in which  $x_n = 1$  in  $M_n$ ,  $n = 1, 2, \dots$ , and  $x_\infty$  in  $C[0, 1]^{**}$  is such that the atomic part of  $x_\infty$  is 0 and the diffuse part of  $x_\infty$  is 1. It is easy to see that for every  $a = (a_n)_{n=1}^\infty$  in  $A$ ,  $(pa^*xp)_n = (pa^*p)_n$ ,  $n = 1, 2, \dots$ , the atomic part of  $(pa^*xp)_\infty$  is 0 and the diffuse part of  $(pa^*xp)_\infty$  is the same as  $(pa^*p)_\infty$ . In particular,  $pa^*xp$  defines a continuous function on  $\overline{X}$  for each  $a$  in  $A$  and so does  $px^*xp$ . But  $x$  does *not* even have a weakly continuous atomic part modulo  $L^{**}$ .

In fact, if there were a  $yp$  in  $\mathcal{W}_p$  such that  $xyp = zyp$ . It follows that the atomic part of  $y_\infty$  is 0. For every continuous function  $f$  in  $C[0, 1]$ , we define an  $a = (a_n)_{n=1}^\infty$  in  $A$  by setting  $a_\infty = f$  in  $C[0, 1]$  and  $a_n = f(h_n)$  in  $M_n$ ,  $n = 1, 2, \dots$ . Since  $yp$  is weakly continuous on  $F(p)$ ,  $pa^*yp = pbp$  for some  $b = (b_n)_{n=1}^\infty$  in  $A$ . In particular,  $b_\infty = 0$  in  $C[0, 1]$  since  $p_\infty = 1$  in  $C[0, 1]^{**}$  and the atomic part of  $a^*y$  is 0. Hence

$$\varphi_n(pa^*yp) = \varphi_n(b) \longrightarrow \varphi_\infty(b) = 0.$$

However,

$$\varphi_n(pa^*yp) = \varphi_n(pa^*xp) = \varphi_n(a^*x) = \varphi_n(a^*)$$

since each  $\varphi_n$  is a pure state in  $F(p)$  and  $x_n = 1$  in  $M_n$ ,  $n = 1, 2, \dots$ . Now, the last term of the equalities approaches to  $\int_0^1 \overline{f(t)} dt$ . Hence,  $\int_0^1 \overline{f(t)} dt = 0$  for every continuous function  $f$  in  $C[0, 1]$  with complex conjugate  $\overline{f(t)}$ , an absurdity! Therefore,  $xp$  does *not* have a weakly continuous atomic part.  $\square$

**Acknowledgement.** We would like to express our deepest gratitude to Professor Lawrence G. Brown for many helpful advice on this work. The current paper as well as [20] is based on the Ph. D. dissertation [19] of the author under his supervision.

#### REFERENCES

- [1] C. A. Akemann, *Left ideal structure of  $C^*$ -algebras*, J. Funct. Anal. **6** (1970), 305–317.
- [2] C. A. Akemann, G. K. Pedersen and J. Tomiyama, *Multipliers of  $C^*$ -algebras*, J. Funct. Anal. **13** (1973), 277–301.
- [3] C. A. Akemann and F. Shultz, *Perfect  $C^*$ -algebras*, Memoirs A. M. S. **326**, 1985.
- [4] C. A. Akemann, J. Andersen and G. K. Pedersen, *Approaching to infinity in  $C^*$ -algebras*, J. Operator Theory **21** (1989), 252–271.
- [5] L. Asimow and A. J. Ellis, *Convexity theory and its applications in functional analysis*, Academic Press, London-New York, 1980.
- [6] L. G. Brown, *Semicontinuity and multipliers of  $C^*$ -algebras*, Can. J. Math. XL (1988), no. 4, 865–988.
- [7] ———, *Complements to various Stone-Weierstrass theorems for  $C^*$ -algebras and a theorem of Shultz*, Commun. Math. Phys. **143** (1992), 405–413.
- [8] ———, *MASA's and certain type I closed faces of  $C^*$ -algebras*, preprint.
- [9] F. Combes, *Quelques propriétés des  $C^*$ -algebras*, Bull. Sci. Math. **94** (1970), 165–192.
- [10] J. Dixmier and A. Douady, *Champs continus d'espaces hilbertiens et de  $C^*$ -algebras*, Bull. Soc. Math. France **91** (1963), 227–284.
- [11] E. G. Effros, *Order ideals in  $C^*$ -algebras and its dual*, Duke Math. **30** (1963), 391–412.

- [12] J. M. G. Fell, *The structure of algebras of operator fields*, Acta Math. **106** (1961), 233–280.
- [13] H. E. Jensen, *Scattered  $C^*$ -algebras*, Math. Scand. **41** (1977), 308–314.
- [14] ———, *Scattered  $C^*$ -algebras, II*, Math. Scand. **43** (1978), 308–310.
- [15] G. K. Pedersen, *Applications of weak\* semicontinuity in  $C^*$ -algebra theory*, Duke Math. J. **39** (1972), 431–450.
- [16] ———,  *$C^*$ -algebras and their automorphism groups*, Academic Press, London, 1979.
- [17] R. T. Prosser, *On the ideal structure of operator algebras*, Memoirs A. M. S. **45**, 1963.
- [18] F. W. Shultz, *Pure states as a dual object for  $C^*$ -algebras*, Commun. Math. Phys. **82** (1982), 497–509.
- [19] Ngai-Ching Wong, *The left quotient of a  $C^*$ -algebra and its representation through a continuous field of Hilbert spaces*, Ph. D. Dissertation, Purdue University, West Lafayette, Indiana, U. S. A., 1991.
- [20] ———, *Left quotients of a  $C^*$ -algebra, I: Representation via vector sections*, J. Operator Theory **32**, 1994, 185–201.

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL SUN YAT-SEN UNIVERSITY, KAOHSIUNG, 80424, TAIWAN, R.O.C.

*E-mail address:* wong@math.nsysu.edu.tw