

Coloring the Cartesian Sum of Graphs

Daphne Der-Fen Liu*

Department of Mathematics
California State University, Los Angeles
Los Angeles, CA 90032, USA
dliu@calstatela.edu

Xuding Zhu†

Department of Applied Mathematics
National Sun Yat-sen University
and
National Center for Theoretical Sciences, Taiwan
zhu@math.nsysu.edu.tw

Submitted May 2005. Revised July 2007 and November 2007.

Abstract

For graphs G and H , let $G \oplus H$ denote their Cartesian sum. We investigate the chromatic number and the circular chromatic number for $G \oplus H$. It is proved that for any graphs G and H , $\chi(G \oplus H) \leq \max\{\lceil \chi_c(G)\chi(H) \rceil, \lceil \chi(G)\chi_c(H) \rceil\}$. It is conjectured that for any graphs G and H , $\chi_c(G \oplus H) \leq \max\{\chi(H)\chi_c(G), \chi(G)\chi_c(H)\}$. We confirm this conjecture for graphs G and H with special values of $\chi_c(G)$ and $\chi_c(H)$. These results improve previously known bounds on the corresponding coloring parameters for the Cartesian sum of graphs.

1991 Mathematics Subject Classification. 05C20, 05C35, 05C15

*Corresponding Author. Supported by the National Science Foundation, USA, under grant DMS 0302456.

†Supported by the National Science Council, Taiwan, under grant NSC93-2115-M-110-004.

Key words and phrases: Cartesian sum of graphs, chromatic number, circular chromatic number, fractional chromatic number.

1 Introduction

Several interesting types of graph products have been studied extensively in the literature. For instance, Klavžar [10] surveyed the study of the chromatic number for four kinds of graph product: Direct product (also known as the categorical product), Cartesian product, lexicographic product and strong product. The circular chromatic number and the fractional chromatic number for these graph products have also attracted considerable attention [13, 17, 18, 19, 22, 24, 25]. Many interesting results are obtained, and yet some challenging problems remain open.

In this article, we investigate the chromatic number and the circular chromatic number for a type of graph products, namely, the Cartesian sum of graphs. Suppose $G = (V, E)$ and $H = (V', E')$. The *Cartesian sum* of G and H , denoted by $G \oplus H$, has as the vertex set $V \times V'$, and the edge set

$$E(G \oplus H) = \{(x, x')(y, y') : xy \in E(G) \text{ or } x'y' \in E(H)\}.$$

This notion of graph product was introduced by Ore [11] in 1962. The chromatic number for the Cartesian sum of graphs has been investigated in [2, 3, 12, 20].

A typical problem on a product of graphs is to express (or find the best bound of) a coloring parameter (such as the chromatic number or the circular chromatic number) for the product of two graphs, say G and H , in terms of the coloring parameters of G and H . For instance, the well-known Hedetniemi conjecture states: For any graph G and H , $\chi(G \times H) = \min\{\chi(G), \chi(H)\}$, where $G \times H$ is the direct product of G and H . Although the conjecture was confirmed for the cases that $\chi(G) = \chi(H) = 3$ by Hedetniemi [7], and $\chi(G) = \chi(H) = 4$ by El-Zahar and Sauer [4], it remains open for other cases.

For any graphs G and H , it is easy to see that $\chi(G \oplus H) \leq \chi(G)\chi(H)$. The bound is sharp, as $\chi(K_n \oplus K_m) = nm$. However, for many graphs, the value of $\chi(G \oplus H)$ is strictly less than $\chi(G)\chi(H)$. Hence, one would aim at finding a better upper bound for $\chi(G \oplus H)$ by investigating further on the structures of G and H . To this end, it is natural to consider the relations between $\chi(G \oplus H)$ and the circular chromatic numbers of G and H .

For a graph G , the circular chromatic number is a refinement of the chromatic number. Let p, q be integers with $p \geq q > 0$. A (p, q) -coloring of a graph G is a mapping $c : V(G) \rightarrow \{0, 1, \dots, p-1\}$ such that for any edge uv of G , $|c(u) - c(v)|_p \geq q$,

where $|x - y|_n = \min\{|x - y|, n - |x - y|\}$. The *circular chromatic number* $\chi_c(G)$ of G is defined as

$$\chi_c(G) = \inf\{p/q : G \text{ admits a } (p, q)\text{-coloring}\}.$$

Notice that the infimum in the above can be replaced by minimum [24]. It is also known [24] that for any graph G ,

$$\chi(G) - 1 < \chi_c(G) \leq \chi(G).$$

The above is equivalent to $\chi(G) = \lceil \chi_c(G) \rceil$. Thus, the circular chromatic number for a graph G contains more information on the structure of G than what the chromatic number does (cf. [24, 25]).

Two questions about the Cartesian sum of graphs emerge naturally:

Question 1 *What is the best upper bound for $\chi(G \oplus H)$ in terms of $\chi_c(G)$ and $\chi_c(H)$?*

Question 2 *What is the best upper bound for $\chi_c(G \oplus H)$ in terms of $\chi_c(G)$ and $\chi_c(H)$?*

As $\chi(G \oplus H) = \lceil \chi_c(G \oplus H) \rceil$, an answer to Question 2 provides an answer to Question 1. However, an answer to Question 1 does not induce an answer to Question 2. In this article, we answer Question 1 completely, and provide partial solutions to Question 2.

A *fractional coloring* of a graph G is a mapping c from $\mathcal{I}(G)$, the set of all independent sets of G , to the interval $[0, 1]$ of reals, such that $\sum_{x \in I \in \mathcal{I}(G)} c(I) \geq 1$ holds for any $x \in V(G)$. The *fractional chromatic number* $\chi_f(G)$ of G is the infimum of the value $\sum_{I \in \mathcal{I}(G)} c(I)$ for a fractional coloring c of G (cf. [5, 8, 14, 15]). It is known [1, 21, 24] that for any graph G , $\chi_f(G) \leq \chi_c(G)$.

For graphs $G = (V, E)$ and $H = (V', E')$, the *lexicographic product* of G and H , denoted by $G[H]$, has the vertex set $V \times V'$, in which $(x, x')(y, y')$ is an edge if $[xy \in E]$ or $[x = y \text{ and } x'y' \in E']$. It was proved in [23] that the following holds for any G and H :

$$\chi_f(G)\chi(H) \leq \chi_c(G[H]) \leq \chi_c(G)\chi(H).$$

If it is the case that $\chi_f(G) = \chi_c(G)$, then we have $\chi_c(G[H]) = \chi_c(G)\chi(H)$. The graphs with $\chi_f(G) = \chi_c(G)$ are called *star-extremal*. Many graphs are known to be star-extremal [6, 9] (for instance, the circular cliques defined later). Hence, there

exist graphs G and H with $\chi_c(G[H]) = \chi_c(G)\chi(H)$. By definition, $G[H]$ is a spanning subgraph of $G \oplus H$. Therefore, if G is star-extremal, then

$$\chi_c(G \oplus H) \geq \chi_c(G)\chi(H).$$

Since the Cartesian sum is symmetric, i.e., $G \oplus H = H \oplus G$, we conclude that if G and H are star-extremal, then

$$\chi_c(G \oplus H) \geq \max\{\chi_c(G)\chi(H), \chi(G)\chi_c(H)\}. \quad (1.1)$$

Consequently, if G and H are star-extremal, then

$$\chi(G \oplus H) \geq \max\{\lceil \chi_c(G)\chi(H) \rceil, \lceil \chi(G)\chi_c(H) \rceil\}. \quad (1.2)$$

To answer Question 1 completely, in Section 2, we prove the following main result:

Theorem 1 *For any graphs G and H ,*

$$\chi(G \oplus H) \leq \max\{\lceil \chi_c(G)\chi(H) \rceil, \lceil \chi(G)\chi_c(H) \rceil\}.$$

Moreover, the bounds are sharp if G and H are star-extremal.

For the circular chromatic number of $G \oplus H$, in Section 3, we prove the following result, giving partial solutions to Question 2:

Theorem 2 *Let p, q, p', q' be positive integers with $p = kq + r$ and $p' = k'q' + r'$, where $0 \leq r < q$ and $0 \leq r' < q'$. Assume $\chi_c(G) = p/q$ and $\chi_c(H) = p'/q'$. Then*

$$\chi_c(G \oplus H) \leq \begin{cases} \chi_c(G)\chi(H), & \text{if } p'(p+q-1) \leq p(p'+q'-r'); \\ \chi_c(H)\chi(G), & \text{if } p(p'+q'-1) \leq p'(p+q-r). \end{cases}$$

Moreover, the bounds are sharp if G and H are star-extremal.

We conjecture that $\chi_c(G \oplus H) \leq \max\{\chi_c(G)\chi(H), \chi(G)\chi_c(H)\}$ holds in general. If the conjecture is proved to be true, then in view of (1.1), this upper bound for $\chi_c(G \oplus H)$ would also be sharp, answering Question 2 completely.

A *homomorphism* of a graph G to another graph H is an edge-preserving mapping from $V(G)$ to $V(H)$. If such a mapping exists, we say that G is *homomorphic* to H , and denote this by $G \rightarrow H$. It is known and easy to see that $\chi(G) \leq n$ if and only if $G \rightarrow K_n$. Two graphs G and H are *homomorphically equivalent*, denoted by

$G \leftrightarrow H$, if $G \rightarrow H$ and $H \rightarrow G$. It is obvious that if $G \leftrightarrow H$, then $\chi(G) = \chi(H)$ and $\chi_c(G) = \chi_c(H)$.

For any given rational $p/q \geq 2$, a *circular clique* $K_{p/q}$ is the graph with the vertex set $V(K_{p/q}) = \{0, 1, 2, \dots, p-1\}$, in which

$$uv \in E(K_{p/q}) \text{ if and only if } |u - v|_p \geq q.$$

In the study of circular chromatic number of graphs, circular cliques play essentially the same role as cliques (complete graphs) in the study of chromatic number. It is known that $\chi_c(K_{p/q}) = \chi_f(K_{p/q}) = p/q$, and for any graph G , $\chi_c(G) \leq p/q$ if and only if $G \rightarrow K_{p/q}$ (cf. [24]). We shall prove that for any p/q and p'/q' , it holds that

$$\chi(K_{p/q} \oplus K_{p'/q'}) = \max\{\lceil (p/q)\lceil p'/q' \rceil \rceil, \lceil (p'/q')\lceil p/q \rceil \rceil\}.$$

This result is used to verify Theorem 1.

Theorem 1 has an application. A useful tool for estimating the chromatic number of a graph $G = (V, E)$ is to partition the edges E into two sets, $E = E_1 \cup E_2$, then determine (or find upper bounds for) the chromatic numbers of the spanning subgraphs $G_1 = (V, E_1)$ and $G_2 = (V, E_2)$. Theorem 1 provides an estimation of $\chi(G)$ when $\chi_c(G_1)$ and $\chi_c(G_2)$ are known.

Corollary 3 *Suppose $G = (V, E)$. Let $E = E_1 \cup E_2$ and $G_i = (V, E_i)$ for $i = 1, 2$. Then*

$$\chi(G) \leq \chi(G_1 \oplus G_2) \leq \max\{\lceil \chi_c(G_1)\chi(G_2) \rceil, \lceil \chi(G_1)\chi_c(G_2) \rceil\}.$$

Assume $G = Q \oplus H$. Let $E_1 = \{(x, x')(y, y') \in E(G) : xy \in E(Q)\}$ and $E_2 = \{(x, x')(y, y') \in E(G) : xy \notin E(Q)\}$, and let $G_1 = (V(G), E_1)$ and $G_2 = (V(G), E_2)$. Then $E(G) = E_1 \cup E_2$, $G_1 \leftrightarrow Q$, and $G_2 \leftrightarrow H$. By Theorem 1, if Q and H are star-extremal, then

$$\begin{aligned} \chi(G) &= \chi(Q \oplus H) = \max\{\lceil \chi_c(Q)\chi(H) \rceil, \lceil \chi(Q)\chi_c(H) \rceil\} \\ &= \max\{\lceil \chi_c(G_1)\chi(G_2) \rceil, \lceil \chi(G_1)\chi_c(G_2) \rceil\}. \end{aligned}$$

This shows that the upper bound in Corollary 3 is sharp.

Note, if it is the case that $\chi_c(G) = \chi(G)$ or $\chi_c(H) = \chi(H)$, then the upper bounds in Theorems 1 and 2 coincide with the trivial bound, $\chi_c(G \oplus H) \leq \chi(G \oplus H) \leq \chi(G)\chi(H)$. Therefore, throughout the article, we assume $\chi_c(G) < \chi(G)$ and $\chi_c(H) < \chi(H)$, when $\chi(G \oplus H)$ or $\chi_c(G \oplus H)$ is in consideration.

2 The Chromatic Number

We determine the chromatic number for the Cartesian sum of circular cliques. Consequences of this result include Theorem 1.

For an integer x and a positive integer m , we denote by $[x]_m$ the unique integer x' such that $0 \leq x' \leq m - 1$, and $x - x'$ is a multiple of m .

Lemma 4 *Let p, q, p', q' be positive integers with $p = kq + r$ and $p' = k'q' + r'$, where $1 \leq r < q$ and $1 \leq r' < q'$. Then*

$$\chi(K_{p/q} \oplus K_{p'/q'}) = \max\{\lceil (p/q)(k' + 1) \rceil, \lceil (p'/q')(k + 1) \rceil\}.$$

Proof. Let $G = K_{p/q} \oplus K_{p'/q'}$. Without loss of generality, we may assume that $(p/q)(k' + 1) \geq (p'/q')(k + 1)$. We shall prove that $\chi(G) = n = \lceil (p/q)(k' + 1) \rceil$.

Since circular cliques are star-extremal, by (1.2), it remains to show that G is n -colorable.

Let $V(G) = \{(i, j) : 0 \leq i \leq p' - 1, 0 \leq j \leq p - 1\}$. For each $(i, j) \in V(G)$, we define the *tile* $B_{i,j}$ by:

$$B_{i,j} = \{([i + a]_{p'}, [j + b]_p) : 0 \leq a \leq q' - 1, 0 \leq b \leq q - 1\}.$$

Intuitively, we regard $V(G)$ as a set of $p \times p'$ grid points embedded on a torus. We draw this torus as a $(p \times p')$ -rectangle R , where the top and the bottom boundaries as well as the left and the right boundaries are identified. As a convention, we assume that the horizontal edge and the vertical edge of R , respectively, has lengths p' and p , as shown in Figure 1.




Figure 1: Covering the torus grid with tiles

Each tile $B_{i,j}$ is a $q \times q'$ rectangle, and covers qq' grid points, with (i, j) as its left-bottom point on R (viewing $(0, 0)$ as the left bottom point of R). By definition of the

Cartesian sum, each $B_{i,j}$ is an independent set of G . To show that G is n -colorable, it suffices to prove that R can be covered by n tiles of the form $B_{i,j}$. Let

$$k'(q-r) = \lambda q + r^*, \text{ where } \lambda, r^* \text{ are integers with } 0 \leq r^* < q.$$

Let

$$z = \begin{cases} \lceil r^*/(q-r) \rceil, & \text{if } r^* < r; \\ 0, & \text{if } r^* \geq r. \end{cases}$$

The n tiles we use to cover R are $B_t = B_{i_t, j_t}, t = 0, 1, \dots, n-1$, where:

$$\begin{aligned} i_t &= \lceil [tq'/(k+1)] \rceil_{p'}, & t &= 0, 1, \dots, n-1; \\ j_t &= \begin{cases} \lceil tq \rceil_p, & \text{if } 0 \leq t < n - z(k+1), \\ \lceil tq - r^* + s(q-r) \rceil_p, & \text{if } n - (s+1)(k+1) \leq t < n - s(k+1), \\ & \text{and } 0 \leq s \leq z-1. \end{cases} \end{aligned}$$

Note that, if $r^* \geq r$, then $z = 0$, and hence $j_t = \lceil tq \rceil_p$ for all t .

Although the above formulas look complicated, the idea about the locations of these tiles is quite simple. Assume $r^* \geq r$ (that is, $z = 0$). We put the first tile, B_0 , at the left-bottom corner of R (see Figure 1). Then we put in the remaining tiles one by one. Assume B_0, B_1, \dots, B_{i-1} are put in (winding around the torus) already. Then we put B_i exactly on the top of B_{i-1} with a ‘‘little bit’’ horizontal shift to the right. The shifts are carefully chosen so that the total of any $k+1$ consecutive shifts is equal to q' . Such a tiling process will wrap around the torus tightly, and as we will show later, the two ends, the last $k+1$ tiles and the first $k+1$ tiles, *meet perfectly* (i.e., without gaps).

Assume $r^* < r$ (that is, $z \geq 1$). We first use the same tiling process in the above, which, for this case, also wraps around the torus tightly, but the two ends may *not* meet perfectly. There might be gaps between the last $k+1$ tiles and the first $k+1$ tiles. To mend this, we divide the last $z(k+1)$ tiles into z ‘‘strips,’’ where each strip consists of consecutive $k+1$ tiles. For instance, the first strip is formed by the last $k+1$ tiles, B_{n-i} for $i = 1, 2, \dots, k+1$, the second strip is formed by the next $k+1$ tiles, $B_{n-(k+1)-i}$ for $i = 1, 2, \dots, k+1$, etc. We shift each of the $(k+1)$ tiles in the first strip vertically downwards by r^* rows, and (if $z \geq 2$) shift each tile in the s -th strip vertically downwards by $r^* - (s-1)(q-r)$ rows, for $2 \leq s \leq z$.

In the following, we claim that these tiles wrap around the torus without gaps. By looking at the first $k+1$ tiles, B_0, B_1, \dots, B_k , we conclude that any horizontal line of R intersects these tiles. In other words, for any $0 \leq b \leq p-1$, there exists some a such that (a, b) is contained in some tile B_t .

Assume that there is a vertex, say (a, b) , which is not covered by any of the tiles. We search, from the point (a, b) on R , along the horizontal line passing through (a, b)

towards the left until we find a vertex which is covered by some tile. To be precise, let $\beta \geq 1$ be the smallest integer such that $([a - \beta]_{p'}, b)$ is contained in some tile. Assume $([a - \beta]_{p'}, b) \in B_t$. As $([a - \beta + 1]_{p'}, b) \notin B_t$, we know that $([a - \beta]_{p'}, b)$ is on the right boundary of B_t , i.e., $[a - \beta]_{p'} = [i_t + q' - 1]_{p'}$. We claim that

$$([a - \beta + 1]_{p'}, b) \in B_{t+k} \cup B_{t+k+1},$$

which is in contrary to the choice of β . (Note, the sub indices of B are taken modulo n .) It suffices to consider two cases.

Case 1. $0 \leq t < n - (k + 1)$ Intuitively, our aim for this case is to show that the tiles, $B_0, B_1, \dots, B_{n-(k+1)}$, wrap around the torus *tightly*, i.e., the right side of each tile B_t , for $t \leq n - (k + 1)$, is covered or touched by the left sides of tiles B_{t+k} and B_{t+k+1} (see Figure 1). To this end, we claim:

$$\{([i_t + x]_{p'}, [j_t + y]_p) : q' \leq x \leq q' + \lfloor kq'/(k+1) \rfloor, 0 \leq y \leq q\} \subseteq B_{t+k} \cup B_{t+k+1}, \quad (2.1)$$

which implies that $([a - \beta + 1]_{p'}, b) \in B_{t+k} \cup B_{t+k+1}$, as $q' \geq 2$.

It follows from the definition that

$$\begin{aligned} [i_{t+k+1} - i_t]_{p'} &= q'; \text{ and} \\ [i_{t+k} - i_t]_{p'} &= \lceil (t+k)q'/(k+1) \rceil - \lceil tq'/(k+1) \rceil. \end{aligned}$$

Hence,

$$\lfloor kq'/(k+1) \rfloor \leq [i_{t+k} - i_t]_{p'} \leq q'.$$

Therefore, to verify (2.1), by noting that the bottom of B_{j+k+1} either touches or overlaps with the top of B_{j+k} , it suffices to show that the following hold:

$$[j_t - j_{t+k}]_p \leq q \quad \text{and} \quad [j_{t+k+1} - j_t]_p \leq q. \quad (2.2)$$

Assume $z = 0$. Since $p = kq + r$, we have

$$\begin{aligned} [j_t - j_{t+k}]_p &= [-kq]_p = r; \\ [j_{t+k+1} - j_t]_p &= [(k+1)q]_p = q - r. \end{aligned}$$

As $r < q$ and $q - r \leq q$, (2.2) follows.

Assume $z \geq 1$. Let $\delta = r^* - (z - 1)(q - r)$. Depending on the values of t , it can be verified that the values of $j_t - j_{t+k}$ and $j_{t+k+1} - j_t$ have the following possibilities:

$$j_t - j_{t+k} \in \{-kq, -(kq - (q - r)), -(kq - \delta)\}$$

and

$$j_{t+k+1} - j_t \in \{(k+1)q, (k+1)q - (q-r), (k+1)q - \delta\}.$$

Again, because $[-kq]_p = r$, $[(k+1)q]_p = q-r$, and $\delta \leq q-r$, we conclude that (2.2) holds for all the above possibilities. This completes the proof of Case 1.

Case 2. $n - (k+1) \leq t < n$ We verify that the last $k+1$ tiles and the first $k+1$ tiles are connected perfectly without any gaps.

Suppose $t = n - (k+1) + m$ for some $0 \leq m \leq k$. First we assume that $r^* < r$. Then, by definition and some calculation, we have $n = (k'+1)k + \lceil (k'+1)r/q \rceil = (k+1)(k'+1) - \lambda$. Therefore,

$$\begin{aligned} j_{n-(k+1)} &= [(n - (k+1))q - r^*]_p \\ &= [(k'(k+1) - \lambda)q - r^*]_p \\ &= [k'(kq + r)]_p \\ &= 0 = j_0. \end{aligned}$$

Thus, for each $0 \leq l \leq k$, we have $j_{n-(k+1)+l} = j_l = lq$. In particular, $j_t = mq$, i.e., B_t and B_m are on the same horizontal level. We shall prove that $([a - \beta + 1]_{p'}, b) \in B_m$. This amounts to show that

$$[i_m - i_{n-(k+1)+m}]_{p'} \leq q'. \quad (2.3)$$

That is,

$$[\lceil mq'/(k+1) \rceil - \lceil (n - (k+1) + m)q'/(k+1) \rceil]_{p'} \leq q'.$$

Since $\lceil mq'/(k+1) \rceil < \lceil (n - (k+1) + m)q'/(k+1) \rceil < p' + \lceil mq'/(k+1) \rceil$, this is equivalent to

$$p' + \lceil mq'/(k+1) \rceil - \lceil (n - (k+1) + m)q'/(k+1) \rceil \leq q'.$$

which holds if

$$p' \leq nq'/(k+1).$$

This is true since $n \geq (p/q)(k'+1) \geq (p'/q')(k+1)$. Notice that by the argument above, (2.3) also holds for $r^* \geq r$.

Next we assume that $r^* \geq r$. Then $n = (k+1)(k'+1) - \lambda - 1$. We shall prove that if $1 \leq m \leq k$, then $([a - \beta + 1]_{p'}, b) \in B_{m-1} \cup B_m$; and if $m = 0$, then $([a - \beta + 1]_{p'}, b) \in B_{n-1} \cup B_0$.

Assume $m \geq 1$. Then $j_{n-(k+1)+m} = \lceil (n - (k+1) + m)q \rceil_p = r^* + (m-1)q$. Hence $j_{m-1} \leq j_t \leq j_m$. To prove that $([a - \beta + 1]_{p'}, b) \in B_{m-1} \cup B_m$, it suffices to show that the right side of the rectangle B_t touches or overlaps with the left sides of B_m

and B_{m-1} . This is equivalent to show that $[i_{m-1} - i_t]_{p'} \leq q'$ and $[i_m - i_t]_{p'} \leq q'$. As $i_t - p' \leq i_{m-1} \leq i_m < i_t$, it suffices to show that $[i_m - i_t]_{p'} = [i_m - i_{n-(k+1)+m}]_{p'} \leq q'$, which is (2.3), and has been verified in the above.

Assume $m = 0$. It amounts to prove that the right side of $B_{n-(k+1)}$ touches or overlaps with the left sides of B_0 and B_{n-1} . This is true since, by a special case of (2.3), we have $[i_0 - i_{n-(k+1)}]_{p'} \leq q'$, and similar to the argument used in Case 1, we have $[i_{n-1} - i_{n-(k+1)}]_{p'} \leq q'$. ■

Proof of Theorem 1) Assume $\chi_c(G) = p/q$ and $\chi_c(H) = p'/q'$. Then there is a homomorphism f from G to $K_{p/q}$, and a homomorphism g from H to $K_{p'/q'}$. It is easy to check that the mapping $\phi(x, y) = (f(x), g(y))$ is a homomorphism from $G \oplus H$ to $K_{p/q} \oplus K_{p'/q'}$. Therefore $\chi(G \oplus H) \leq \chi(K_{p/q} \oplus K_{p'/q'})$. The conclusion then follows from Lemma 4. The moreover part follows from (1.2). ■

The next result follows from Theorem 1 and (1.2).

Corollary 5 *For any two graphs G and H , if $\chi_c(G)\chi(H) \geq \chi(G)\chi_c(H)$ and G is star-extremal, then*

$$\chi(G \oplus H) = \lceil \chi_c(G)\chi(H) \rceil.$$

Čížek and Klavžar [3] showed that if G and H are vertex-critical and are not complete graphs, then $\chi(G \oplus H) < \chi(G)\chi(H)$. It is known [16] that if G is vertex-critical with $\chi(G) = k$ and with girth (length of a shortest cycle) at least k , then $\chi_c(G) \leq k - (1/2)$. Hence, by Theorem 1, we have

Corollary 6 *Let G be a vertex-critical graph with $\chi(G) = k$ and girth at least k . If H is a graph such that $\chi_c(G)\chi(H) \geq \chi(G)\chi_c(H)$, then*

$$\chi(G \oplus H) \leq \lceil (k - 1/2)\chi(H) \rceil.$$

3 The Circular Chromatic Number

Theorem 1 gives the best upper bound for the chromatic number of the Cartesian sum $G \oplus H$ in terms of $\chi_c(G)$ and $\chi_c(H)$. However, the best upper bound for the circular chromatic number of $G \oplus H$ in terms of $\chi_c(G)$ and $\chi_c(H)$ for all graphs G and H remains unknown. We propose:

Conjecture 1 *For any graphs G and H ,*

$$\chi_c(G \oplus H) \leq \max\{\chi_c(G)\chi(H), \chi(G)\chi_c(H)\}.$$

As we pointed out earlier, if Conjecture 1 is true, then this upper bound would be the best. Theorem 2 confirms the conjecture for some special graphs.

To prove Theorem 2, we make use of the following result.

Lemma 7 *Let p, q, p', q' be positive integers with $p = kq + r$ and $p' = k'q' + r'$, where $1 \leq r < q$ and $1 \leq r' < q'$. Then*

$$\chi_c(K_{p'/q'} \oplus K_{p/q}) = \begin{cases} (p/q)(k' + 1), & \text{if } p'(p + q - 1) \leq p(p' + q' - r'); \\ (p'/q')(k + 1), & \text{if } p(p' + q' - 1) \leq p'(p + q - r). \end{cases}$$

Proof. By (1.1), $\chi_c(K_{p/q} \oplus K_{p',q'}) \geq \max\{(p/q)(k' + 1), (p'/q')(k + 1)\}$. Thus, we only need to verify the other direction of the inequality.

By symmetry, it is enough to verify the second case in Lemma 7. Assume $p(p' + q' - 1) \leq p'(p + q - r)$. Observe that $(p'/q')(k + 1) \geq (p/q)(k' + 1)$ is equivalent to $p(p' + q' - 1) \leq p'(p + q - r)$. So $p(p' + q' - 1) \leq p'(p + q - r)$ implies that

$$(p'/q')(k + 1) \geq (p/q)(k' + 1).$$

Let $G = K_{p'/q'} \oplus K_{p/q}$. It suffices to find an (n, q') -coloring for G , where $n = p'(k + 1)$. Before defining such a coloring, we need a few more notations. Divide $p' + q' - 1$ into q numbers *almost evenly*, by letting

$$t_i = \lceil i(p' + q' - 1)/q \rceil - \lceil (i - 1)(p' + q' - 1)/q \rceil, \quad i = 1, 2, \dots, q.$$

Then, $t_1 + t_2 + \dots + t_l = \lceil l(p' + q' - 1)/q \rceil$ for any $1 \leq l \leq q$.

Define y_j for $j = 0, 1, \dots, p - 1$ by:

$$\begin{aligned} y_0 &= 0; \text{ and} \\ y_j &= y_{j-1} + t_b, \text{ if } j \geq 1 \text{ and } j = sq + b, \text{ where } s, b \text{ are integers and } 1 \leq b \leq q. \end{aligned}$$

For any integers z, w , the *interval from z to w modular n* , denoted by $[z, w]_n$, is the set of consecutive integers from $[z]_n$ to $[w]_n$, with calculation taken under modulo n . For instance, $[2, 5]_7 = \{2, 3, 4, 5\}$, $[5, 2]_7 = \{5, 6, 0, 1, 2\}$ and $[13, 8]_7 = \{6, 0, 1\}$.

For each $0 \leq j \leq p - 1$, let I_j be the interval:

$$I_j = [y_j, y_j + p' - 1]_n.$$

For $0 \leq x \leq n - 1$, let:

$$A_x = \{(i, j) : i = [x]_{p'} \text{ and } x \in I_j\}.$$

Notice that $A_x \cap A_y = \emptyset$ if $x \neq y$. For if $(i, j) \in A_x \cap A_y$ for some $x \neq y$, then by definition, $|x - y| = mp'$ for some positive integer m , and $x, y \in I_j$. This is impossible, as I_j consists of p' consecutive integers.

The coloring f is defined by $f(i, j) = x$, if $(i, j) \in A_x$. It remains to show that f is an (n, q') -coloring for G . Suppose $(i, j)(i', j')$ is an edge in G . Let $f(i, j) = x$ and $f(i', j') = y$. We need to prove that $|x - y|_n \geq q'$.

By definition, $ii' \in E(K_{p', q'})$ or $jj' \in E(K_{p, q})$. If $ii' \in E(K_{p', q'})$, then $|i - i'|_{p'} \geq q'$. By definition, $i = [x]_{p'}$ and $i' = [y]_{p'}$, implying $|x - y|_n \geq q'$ (since $n = p'(k + 1)$).

Assume $jj' \in E(K_{p, q})$. By definition, $x \in I_j$, $y \in I_{j'}$ and $|j - j'|_p \geq q$. Hence,

$$y \in I_{j+q} \cup I_{j+q+1} \cup \cdots \cup I_{j+p-q},$$

with the sub-indices taken under modulo p . This implies,

$$y \in [y_{j+q}, y_{j+p-q} + p' - 1]_n.$$

Let $j = sq + b$ for some $1 \leq b \leq q$. Then, we have

$$x \in I_j = [y_j, y_j + p' - 1]_n,$$

where $y_j = s(p' + q' - 1) + t_1 + t_2 + \cdots + t_b$.

Hence, to show that $|x - y|_n \geq q'$, it suffices to verify the following:

$$q' \leq |y_{j+q} - (y_j + p' - 1)| \leq n - q' \tag{3.1}$$

$$q' \leq |(y_{j+p-q} + p' - 1) - y_j| \leq n - q'. \tag{3.2}$$

Since $y_{j+q} = y_j + p' + q' - 1$, so $|y_{j+q} - (y_j + p' - 1)| = q'$. Thus, (3.1) holds. As $j + p - q = j + (k - 1)q + r$, we have

$$y_{j+p-q} = y_j + (k - 1)(p' + q' - 1) + t_{b+1} + t_{b+2} + \cdots + t_{b+r}, \text{ implying}$$

$$(y_{j+p-q} + p' - 1) - y_j = k(p' + q' - 1) + t_{b+1} + \cdots + t_{b+r} - q'.$$

Since $k \geq 2$, we have $(y_{j+p-q} + p' - 1) - y_j \geq q'$. It remains to show that $(y_{j+p-q} + p' - 1) - y_j \leq n - q'$. As $t_{b+1} + \cdots + t_{b+r} \leq \lceil (r/q)(p' + q' - 1) \rceil$, to prove (3.2), it suffices to show that

$$k(p' + q' - 1) + \lceil (r/q)(p' + q' - 1) \rceil \leq n.$$

which is equivalent to

$$k(p' + q' - 1) + (r/q)(p' + q' - 1) \leq n.$$

The above is true by the assumption that $p(p' + q' - 1) \leq p'(p + q - r)$. This completes the proof. ■

Proof of Theorem 2) Suppose $\chi_c(G) = p/q$ and $\chi_c(H) = p'/q'$. Then G admits a homomorphism to $K_{p/q}$, and H admits a homomorphism to $K_{p'/q'}$, which implies that $G \oplus H$ admits a homomorphism to $K_{p/q} \oplus K_{p'/q'}$. Therefore,

$$\chi_c(G \oplus H) \leq \chi_c(K_{p/q} \oplus K_{p'/q'}).$$

The conclusion of Theorem 2 then follows from Lemma 7. The moreover part follows from (1.1). ■

Corollary 8 *If $\chi_c(G) = p/q$, $[p]_q = 1$ and $\chi_c(G)\chi(H) \geq \chi(G)\chi_c(H)$ then*

$$\chi_c(G \oplus H) \leq \chi_c(G)\chi(H).$$

Moreover, the equality holds if G is star-extremal.

Proof. Assume $p = kq + 1$ and $\chi_c(H) = p'/q'$ where $p' = k'q' + r'$. The hypothesis $\chi_c(G)\chi(H) \geq \chi(G)\chi_c(H)$ is equivalent to $p'(p + q - 1) \leq p(p' + q' - r')$. The moreover part follows from (1.1). ■

Another consequence of Theorem 7 is the case when $r = r' = 1$, for which, by Corollary 8, the value of $\chi_c(K_{kq+1} \oplus K_{k'q'+1})$ can be completely determined, confirming Conjecture 1 for this case.

Corollary 9 *For any graphs G and H with $\chi_c(G) = p/q$ and $\chi_c(H) = p'/q'$. If $[p]_q = 1$ and $[p']_{q'} = 1$, then*

$$\chi_c(G \oplus H) \leq \max\{\chi_c(G)\chi(H), \chi(G)\chi_c(H)\}.$$

Moreover, the equality holds if G and H are star-extremal.

In searching of the graphs with $\chi(G \oplus H) < \chi(G)\chi(H)$, Borowiecki [2] pointed out that $\chi(C_5 \oplus C_5) \leq 8$. By Corollary 9 and the fact that $\chi_c(C_{2k+1}) = \chi_f(C_{2k+1}) = 2 + (1/k)$ for odd cycles, a stronger and more general result emerges:

Example 1 *For any $n \geq k \geq 2$, $\chi_c(C_{2n+1} \oplus C_{2k+1}) = 6 + (3/k)$.*

Acknowledgment

The authors thank Claude Tardif for his comments on a problem related to the chromatic number of the Cartesian sum of odd cycles, which motivated the study of this work. They also wish to thank the two anonymous referees for constructive comments which helped to improve the presentation of this article. The coloring used in the proof of Theorem 1 was initially inspired by several examples of “covering the dots on a donut” by Jessica Ma.

References

- [1] J. A. Bondy and P. Hell, *A note on the star chromatic number*, J. Graph Theory, 14 (1990), 479 – 482.
- [2] M. Borowiecki, *On chromatic number of products of two graphs*, Colloq. Math., 25 (1972), 49 – 52.
- [3] N. Čížek and S. Klavžar, *On the chromatic number of the lexicographic product and the Cartesian sum of graphs*, Discrete Math., 134 (1994), 17 – 24.
- [4] M. El-Zahar and N. Sauer, *The chromatic number of the product of two 4-chromatic graphs is 4*, Combinatorica, 5 (1985), 121 – 126.
- [5] D. Fisher, *Fractional coloring with large denominators*, J. Graph Theory, 20 (1995), 403 – 409.
- [6] G. Gao and X. Zhu, *Star-extremal graphs and the lexicographic product*, Disc. Math., 152 (1996), 147 – 156.
- [7] S. T. Hedetniemi, *Homomorphisms of graphs and automata*, Technical Report 03105–44–T, University of Michigan, 1966.
- [8] I. Leader, *The fractional chromatic number of infinite graphs*, J. Graph Theory, 20 (1995), 411 – 417.
- [9] K.-W. Lih, D. Liu and X. Zhu, *Star-extremal circulant graphs*, SIAM J. Disc. Math., 12 (1999), 491 – 499.
- [10] S. Klavžar, *Coloring graph products - A survey*, Disc. Math., 155 (1996), 135—145.
- [11] O. Ore, *Theory of Graphs*, American Mathematical Society, 1962.
- [12] V. Puš, *Chromatic number of products of graphs*, Technical Report 88-85, Charles University, Prague, 1988.

- [13] N. Sauer, *Hedetniemi's conjecture – a survey*, Discrete Math., 229 (2001), 261 – 292.
- [14] E. Scheinerman and D. Ullman, *Fractional Graph Theory*, Wiley-Interscience Series in Discrete Mathematics and Optimization, 1997.
- [15] S. H. Scott, *Multiple node colourings of finite graphs*, Doctoral dissertation, University of Reading, England, March 1975.
- [16] E. Steffen and X. Zhu, *On the star chromatic numbers of graphs*, Combinatorica, 16 (1996), 439 – 448.
- [17] C. Tardif, *The chromatic number of the product of two graphs is at least half the minimum of the fractional chromatic numbers of the factors*, Comment. Math. Univ. Carolin., 42 (2001), 353 – 355.
- [18] C. Tardif, *Chromatic numbers of products of tournaments: fractional aspects of Hedetniemi's conjecture*, Graphs, homomorphism and statistical physics, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 63 (2004), 171 – 175.
- [19] C. Tardif, *Multiplicative graphs and semi-lattice endomorphisms in the category of graphs*, J. Combin. Theory, Ser. B, 95 (2005), no. 2, 338–345.
- [20] K. W. Yang, *Chromatic number of Cartesian sum of two graphs*, Proc. Amer. Math. Soc., 19 (1968), 617 – 618.
- [21] A. Vince, *Star chromatic number*, J. Graph Theory, 12 (1988), 551 – 559.
- [22] X. Zhu, *The fractional chromatic number of the direct product of graphs*, Glasgow Mathematical Journal, 44 (2002), 103 – 115.
- [23] X. Zhu, *Star chromatic numbers and products of graphs*, J. Graph Theory, 16 (1992), 557 – 569.
- [24] X. Zhu, *Circular chromatic number: A survey*, Disc. Math., 229 (2001), 371 – 410.
- [25] X. Zhu, *Recent developments in circular colouring of graphs*. Topics in discrete mathematics, 497–550, Algorithms Combin., 26, Springer, Berlin, 2006.
- [26] X. Zhu, *A survey on Hedetniemi's conjecture*, Taiwanese J. Math., 2 (1998), 1 – 24.