

# Density of the circular chromatic numbers of series-parallel graphs

Zhishi Pan and Xuding Zhu\*  
Department of Applied Mathematics  
National Sun Yat-sen University  
Kaohsiung, Taiwan 80424  
Email: zhu@math.nsysu.edu.tw

## Abstract

Suppose  $G$  is a series-parallel graph. It was proved in [3] that either  $\chi_c(G) = 3$  or  $\chi_c(G) \leq 8/3$ . So none of the rationals in the interval  $(8/3, 3)$  is the circular chromatic number of a series-parallel graph. This paper proves that for every rational  $r \in [2, 8/3] \cup \{3\}$  there exists a series-parallel graph  $G$  with  $\chi_c(G) = r$ .

## 1 Introduction

The circular chromatic number (also known as the star chromatic number)  $\chi_c(G)$  of a graph  $G$  is a natural generalization of the chromatic number of a graph. There are quite a few equivalent definitions for the circular chromatic number of a graph [10, 11, 14]. In this paper we use the following definition:

Suppose  $r \geq 1$  is a real number. An  $r$ -colouring of a graph  $G$  is a mapping  $f : V(G) \rightarrow [0, r)$  such that for every edge  $uv$  of  $G$  we have  $1 \leq |f(u) - f(v)| \leq r - 1$ . We say  $G$  is  $r$ -colourable if there exists an  $r$ -colouring of  $G$ . In case  $r = k$  is an integer, then  $r$ -colourability coincides with the ordinary vertex  $k$ -colourability. The *circular chromatic number*  $\chi_c(G)$  of  $G$  is the infimum of those  $r$  for which  $G$  is  $r$ -colourable.

---

\*This research was partially supported by the National Science Council under grant NSC 89-2115-M-110-003

It is known [14] that for a finite graph  $G$ , the infimum in the definition is always attained and hence can be replaced by the minimum. Moreover for a finite graph  $G$ , the circular chromatic number of  $G$  is always a rational.

If  $r' \geq r$  and  $G$  is  $r$ -colourable then  $G$  is  $r'$ -colourable. Since for an integer  $r = k$ ,  $r$ -colourability coincides with the ordinary  $k$ -colourability, so for any finite graph  $G$  we have

$$\chi(G) - 1 < \chi_c(G) \leq \chi(G).$$

Therefore  $\chi(G) = \lceil \chi_c(G) \rceil$ . So the circular chromatic number  $\chi_c(G)$  is a refinement of the chromatic number  $\chi(G)$ , and  $\chi(G)$  is an approximation of  $\chi_c(G)$ .

It follows from the definition that any non-trivial graph has circular chromatic number at least 2. On the other hand, it was shown in [1, 10] that for any rational number  $r \geq 2$ , there exists a finite graph  $G$  of circular chromatic number  $r$ .

However if we restrict to graphs with a certain property, then the values for their circular chromatic numbers maybe restricted. The well-known Hadwiger's conjecture says that  $K_n$ -minor free graphs have chromatic number at most  $n - 1$ . If this conjecture is true then the circular chromatic numbers of  $K_n$ -minor free graphs are also at most  $n - 1$ . An interesting question related to this conjecture is this:

**Question 1.1** *Is it true that for every rational number  $2 \leq r \leq n - 1$  there is a  $K_n$ -minor free graph  $G$  with  $\chi_c(G) = r$  ?*

The question has been studied in [3, 5, 12, 13, 4], and now we have a complete solution. It is proved in [4] that every rational  $r \in [2, n - 1]$  is the circular chromatic number of a  $K_n$ -minor free graph if  $n \geq 5$ . In sharp contrast to this, the answer is negative for  $n = 4$ . The following result is proved in [3]:

**Theorem 1.1** *If  $G$  is a  $K_4$ -minor free graph then either  $\chi_c(G) = 3$  or  $\chi_c(G) \leq 8/3$ .*

So the interval  $(8/3, 3)$  is a gap among the circular chromatic numbers of  $K_4$ -minor free graphs. A natural question is: are there other gaps among the circular chromatic numbers of  $K_4$ -minor free graphs ?

This paper answers this question: there are no other gaps. To be precise we shall prove the following result:

**Theorem 1.2** *Suppose  $r$  is a rational number. If  $r = 3$  or  $2 \leq r \leq 8/3$  then there exists a  $K_4$ -minor free graph  $G$  with  $\chi_c(G) = r$ .*

## 2 Two terminal series-parallel graphs

The class of  $K_4$ -minor free graphs can be defined in many different ways and is referred to by different names, such as series-parallel graphs, partial 2-trees, etc., [3, 7]. We adopt the following definition of two-terminal series-parallel graphs from [3]. A two-terminal series-parallel graph  $(G; x, y)$  is defined recursively as follows:

- Let  $V(K_2) = \{0, 1\}$ . Then  $(K_2; 0, 1)$  is a two-terminal series-parallel graph.
- (The parallel construction.) Let  $(G; x, y)$  and  $(G'; x', y')$  be two disjoint two-terminal series-parallel graphs. Define  $G''$  to be the graph obtained from the union of  $G$  and  $G'$  by identifying  $x$  and  $x'$  into a single vertex  $x''$ , and identifying  $y$  and  $y'$  into a single vertex  $y''$ . Then  $(G''; x'', y'')$  is a two-terminal series-parallel graph.
- (The series construction.) Let again  $(G; x, y)$  and  $(G'; x', y')$  be two disjoint two-terminal series-parallel graphs. Define  $G''$  to be the graph obtained from the union of  $G$  and  $G'$  by identifying  $y$  and  $x'$  into a single vertex. Then  $(G''; x, y')$  is a two-terminal series-parallel graph.
- There are no other two-terminal series-parallel graphs.

A graph  $G$  is a series-parallel graph if there exist some two vertices  $x, y$  such that  $(G; x, y)$  is a two-terminal series-parallel graph. For all the series-parallel graphs in the remaining part, there are always two terminals which are clearly indicated in the context. Moreover, if the series-parallel graph is denoted by  $G$  (resp.  $G', G''$  etc.) then the two terminals are denoted by  $x$  and  $y$  (resp.  $x'$  and  $y'$ ,  $x''$  and  $y''$ , etc.). The circular chromatic number of series-parallel graphs has been studied in [2, 3, 8, 9].

It is well-known [7] that a graph  $G$  is  $K_4$ -minor free if and only if every block of  $G$  is a series-parallel graph. Therefore it suffices to prove Theorem 1.2 for series-parallel graphs, i.e., we shall prove the following:

**Theorem 2.1** *Suppose  $r$  is a rational number. If  $r = 3$  or  $2 \leq r \leq 8/3$  then there is a series-parallel graph  $G$  with  $\chi_c(G) = r$ .*

### 3 The labeling method

For a real number  $r \geq 1$ , an  $r$ -colouring of a graph  $G$  uses all the real numbers in the interval  $[0, r)$  as colours. We view the numbers in  $[0, r)$  as cyclically ordered, i.e., we identify the two ends 0 and  $r$  into a single point to obtain a circle  $C^r$  of circumference  $r$ , as depicted in Fig. 1 below. Thus an  $r$ -colouring  $f$  of  $G$  “colours” the vertices of  $G$  with points of  $C^r$ , and the points assigned to adjacent vertices must be at least unit distance apart.

As a graph with at least one edge has circular chromatic number at least 2, in the following we shall only consider  $r$ -colouring for  $r \geq 2$ .




Figure 1: Illustration of the circle  $C^r$  for  $r = 3.5$  and intervals of  $C^r$

Suppose  $r \geq 2$  is a real number. For any two real numbers  $a, b \in [0, r)$ , we denote by  $[a, b]_r$  the interval of  $C^r$  goes from  $a$  to  $b$  along the clockwise (i.e., the increasing) direction. To be precise, if  $a < b$  then the interval  $[a, b]_r$  contains all the real numbers  $t$  such that  $a \leq t \leq b$ ; if  $a > b$  then the interval  $[a, b]_r$  contains all the real numbers  $t$  such that either  $a \leq t < r$  or  $0 \leq t \leq b$ . The case  $a = b$  is ambiguous (it could be the whole circle  $C^r$  or just a single point). This case rarely occurs in this paper and for the few places it occurs, the meaning will be clear from the context. When the real number  $r$  is clear from the context, we shall write  $[a, b]$  for  $[a, b]_r$ .

For  $r = 3.5$ , the intervals  $[0.5, 2]$  and  $[2.5, 0.2]$  are depicted in Fig. 1. The length  $\ell([a, b]_r)$  of an interval  $[a, b]_r$  of  $C^r$  is just the geometric length of that interval on  $C^r$ , i.e., if  $a < b$  then  $\ell([a, b]_r) = b - a$ ; if  $a > b$  then

$$\ell([a, b]_r) = b + r - a.$$

Given a series-parallel graph  $G$  with terminals  $x$  and  $y$ . The  $r$ -label set  $L_r(G)$  of  $G$  is a subset of  $C^r$  defined as follows:

$$L_r(G) = \{t \in C^r : \text{there is an } r\text{-colouring } f \text{ of } G \text{ such that } f(x) = 0 \text{ and } f(y) = t\}.$$

It follows from this definition that  $G$  is  $r$ -colourable if and only if  $L_r(G) \neq \emptyset$ .

For two subsets  $A, B$  of  $C^r$ , we define

$$A + B = \{t : \exists a \in A, b \in B, a + b \bmod r = t\}.$$

**Lemma 3.1** *Suppose  $r \geq 2$  is a real number. The  $r$ -label set  $L_r(G)$  of all series-parallel graphs  $G$  can be calculated recursively as follows:*

1. *If  $G = K_2$  then  $L_r(G) = [1, r - 1]$ ;*
2. *If  $G$  is obtained from  $G_1$  and  $G_2$  by a parallel construction then  $L_r(G) = L_r(G_1) \cap L_r(G_2)$ ;*
3. *If  $G$  is obtained from  $G_1$  and  $G_2$  by a series construction then  $L_r(G) = L_r(G_1) + L_r(G_2)$ .*

The proof of Lemma 3.1 is straightforward and omitted.

**Lemma 3.2** *Suppose  $A = [a, b]_r$  and  $B = [c, d]_r$ . If  $\ell([a, b]_r) + \ell([c, d]_r) \geq r$  then  $A + B = C^r$ ; if  $\ell([a, b]_r) + \ell([c, d]_r) < r$  then  $A + B = [a + c, b + d]$ , where the summations are carried out modulo  $r$ .*

The proof of Lemma 3.2 is also straightforward and omitted.

Note that if  $A = \emptyset$  then for any set  $B$ ,  $A + B = \emptyset$ .

It is well-known that series-parallel graphs have circular chromatic numbers at most 3. Moreover, the only series-parallel graphs with circular chromatic number 3 are those that contain triangles (see [3]), which we are not interested in this paper. So in the remaining of this paper, we assume that  $2 \leq r < 3$ . Denote by  $P_i$  the path of length  $i$  (i.e., the path with  $i$  edges).

**Example 3.1** *Suppose  $2 \leq r < 3$ . Let  $C_5$  be the pentagon, and let  $G$  be the graph as depicted in Fig. 2. Then*

- $L_r(P_2) = [2, r - 2]$ .
- $L_r(P_3) = [3 - r, 2r - 3]$ .
- If  $r < 5/2$  then  $L_r(C_5) = \emptyset$ , if  $5/2 \leq r < 3$  then  $L_r(C_5) = [2, 2r - 3] \cup [3 - r, r - 2]$ .
- If  $r < 8/3$  then  $L_r(G) = \emptyset$ , if  $8/3 \leq r < 3$  then  $L_r(G) = [2, 3r - 6] \cup [5 - r, 2r - 3] \cap [3 - r, 2r - 5] \cup [6 - 2r, r - 2]$ .




Figure 2: The graph  $G$  and  $G'$  for Example 3.1

**Proof.** The path  $P_2$  is obtained from two copies of  $K_2$  by a series construction. By Lemma 3.1,

$$L_r(P_2) = L_r(K_2) + L_2(K_2) = [1, r - 1] + [1, r - 1] = [2, r - 2].$$

The path  $P_3$  is obtained from  $P_2$  and  $K_2$  by a series construction. So

$$L_r(P_3) = L_r(P_2) + L_r(K_2) = [3, 2r - 3].$$

The graph  $C_5$  is obtained from  $P_2$  and  $P_3$  by a parallel construction. So

$$L_r(C_5) = L_r(P_2) \cap L_r(P_3).$$

Thus if  $r < 5/2$  then  $L_r(C_5) = \emptyset$  and if  $r \geq 5/2$  then

$$L_r(C_5) = [2, 2r - 3] \cup [3 - r, r - 2].$$

Now to construct the graph  $G$  as depicted in Fig. 2, we first do a series construction with two copies of  $C_5$  to obtain the graph  $G'$  as depicted in Fig.

2, then do a parallel construction with  $C_5$  and  $G'$ . By using Lemmas 3.1 and 3.2, when  $r \geq 5/2$  we have

$$\begin{aligned} L_r(G') &= L_r(C_5) + L_r(C_5) \\ &= [4 - r, 3r - 6] \cup [6 - 2r, 2r - 4] \cup [5 - r, 2r - 5]. \end{aligned}$$

Now if  $r < 8/3$ , then  $L_r(G) = L_r(G') \cap L_r(C_5) = \emptyset$ , if  $r \geq 8/3$  then

$$\begin{aligned} L_r(G) &= L_r(G') \cap L_r(C_5) \\ &= [2, 3r - 6] \cup [5 - r, 2r - 3] \cap [3 - r, 2r - 5] \cup [6 - 2r, r - 2]. \end{aligned}$$

■

It follows from the discussion above that  $\chi_c(C_5) = 5/2$  and  $\chi_c(G) = 8/3$  for the graph  $G$  as depicted in Fig. 2.

## 4 Proof of Theorem 2.1

The triangle  $K_3$  is a series-parallel graph with  $\chi_c(K_3) = 3$ ,  $K_2$  is a series-parallel graph with  $\chi_c(K_2) = 2$ , and the example in Section 3 shows that there is a series-parallel graph  $G$  with  $\chi_c(G) = 8/3$ . In this section, for each fraction  $p/q \in (2, 8/3)$  we shall construct a series-parallel graph  $G$  with  $\chi_c(G) = p/q$ .

The construction of these graphs is by induction on the denominator  $q$ . For each reduced fraction  $p/q \in (2, 3)$  we define the parents of  $p/q$  as follows: We construct a table in the following way. In the first row we write  $2/1$  and  $3/1$ . For  $n = 2, 3, \dots$  we use the rule: Form the  $n$ th row by copying the  $(n - 1)$ st in order, but insert the fraction  $(a + a')/(b + b')$  between the consecutive fractions  $a/b$  and  $a'/b'$  if  $b + b' \leq n$ . For example, the second row is

$$2/1, 5/2, 3/1,$$

the third row is

$$2/1, 7/3, 5/2, 8/3, 3/1,$$

the fourth row is

$$2/1, 9/4, 7/3, 5/2, 8/3, 11/4, 3/1.$$

and the fifth row is

$$2/1, 11/5, 9/4, 7/3, 12/5, 5/2, 13/5, 8/3, 11/4, 14/5, 3/1.$$

Suppose the fraction  $p/q$  is constructed in the process above as  $p/q = (a + a')/(b + b')$ . We shall call the fraction  $a/b, a'/b'$  the *lower parent* and the *upper parent* of  $p/q$ , respectively. We denote the lower parent of  $p/q$  by  $p_l(p/q)$  and denote the upper parent of  $p/q$  by  $p_u(p/q)$ .

The  $n$ th row of the table constructed above is called the *Farey sequence of order  $n$*  [6]. It is well-known (see page 298 of [6]) that the  $n$ th row contains all the fractions in the set  $[2, 3]$  with denominator not exceeding  $n$ , and if  $a/b$  and  $a'/b'$  are two consecutive fractions of the  $n$ th row then  $a'b - ab' = 1$ . So each of the fractions  $p/q \in [2, 3]$  has a unique lower parent and a unique upper parent, except for  $2/1$  and  $3/1$  which have no parents.

**Lemma 4.1** *Suppose  $a/b < a'/b'$  are two consecutive fractions of the  $n$ th row for some  $n$ . If  $b < b'$ , then  $(a' - 2)/(b' - 1) \leq (a - 2)/(b - 1)$ . Moreover,  $(a + a' - 2)/(b + b' - 1) \leq (a' - 2)/(b' - 1)$ .*

**Proof.** Since  $a/b < a'/b'$  and  $b < b'$  we have

$$b'(a'/b' - 2) - b(a/b - 2) > 0.$$

So

$$b'(a'/b' - 2) - b(a/b - 2) = a' - 2b' - a + 2b \geq 1.$$

As  $a'b - ab' = 1$ , we have

$$a' - 2b' - a + 2b \geq a'b - ab'.$$

Therefore

$$(a' - 2)(b - 1) = a'b - a' - 2b + 2 \leq ab' - 2b' - a + 2 = (a - 2)(b' - 1),$$

i.e.,

$$(a' - 2)/(b' - 1) \leq (a - 2)/(b - 1).$$

For the moreover part, since  $a/b > 2$  we have

$$2b - a < 0 < a'b - ab' = 1.$$

Therefore  $ab' - a \leq a'b - 2b$ , which implies that

$$ab' + a'b' - 2b' - a - a' + 2 \leq a'b + a'b' - a' - 2b - 2b' + 2,$$

i.e.,

$$(a + a' - 2)/(b + b' - 1) \leq (a' - 2)/(b' - 1).$$

■

**Corollary 4.1** *If  $p_l(p/q) = a/b$  and  $p_u(p/q) = a'/b'$  then  $(p-2)/(q-1) \leq (a-2)/(b-1)$  and  $(p-2)/(q-1) \leq (a'-2)/(b'-1)$ .*

**Proof.** If  $p_l(p/q) = a/b$  then  $a/b < p/q$  are two consecutive fractions of the  $n$ th row for some  $n$  and  $b < q$ . The second half follows from the moreover part of Lemma 4.1. ■

**Lemma 4.2** *Suppose  $p/q \in (2, 8/3]$ ,  $p_l(p/q) = a/b$  and  $p_u(p/q) = a'/b'$ . Then there is a series-parallel graph  $G_{p/q}$  such that if  $a/b \leq r < \min\{8/3, (p-2)/(q-1)\}$  then*

$$L_r(G_{p/q}) = [p-3-(q-2)r, (q-1)r-p+3],$$

if  $r < a/b$  then

$$L_r(G_{p/q}) = \emptyset.$$

**Proof.** We shall prove Lemma 4.2 by induction on  $q$ . If  $q = 2$ , i.e.,  $p/q = 5/2$ , then  $a/b = 2/1$ . Let  $G_{5/2} = P_2$ . Then  $L_r(G_2) = [2, r-2]$  for  $2 \leq r < 8/3$ . If  $p/q = 8/3$  then  $a/b = 5/2$ . Let  $G_{8/3}$  be the graph as depicted in Fig. 3. It is straightforward to verify (by using Lemmas 3.1 and 3.2) that for  $5/2 \leq r < 8/3$   $L_r(G_4) = [5-r, 2r-5]$ .




Figure 3: The graph  $G_{8/3}$

Suppose  $q \geq 3$  and  $p/q \neq 8/3$ , and Lemma 4.2 is true for all the fractions in  $p'/q'(2, 8/3]$  with  $q' < q$ .

First we consider the case that  $a/b = 2/1$ . Then  $p/q = (2h+1)/h$  for some integer  $h \geq 3$ . Let  $G_{(2h+1)/h} = P_{2h-2}$ . By Lemma 3.1 and Lemma 3.2, it is straightforward to verify that  $L_r(G_{(2h+1)/h}) = \emptyset$  for  $r < 2$  and  $L_r(G_{(2h+1)/h}) = [2h-2-(h-2)r, (h-1)r-2h+2]$  for  $2 \leq r < (2h-1)/(h-1)$ .

In the remaining we assume that  $p_l(p/q) = a/b, p_u(p/q) = a'/b' \in (2, 8/3]$ . By the induction hypothesis, there is a graph  $G_{a/b}$  such that if  $p_l(a/b) \leq r < (a-2)/(b-1)$  then

$$L_r(G_{a/b}) = [a-3-(b-2)r, (b-1)r-a+3],$$

if  $r < p_l(a/b)$  then

$$L_r(G_{a/b}) = \emptyset.$$

There is a graph  $G_{a'/b'}$  such that if  $p_l(a'/b') \leq r < (a' - 2)/(b' - 1)$  then

$$L_r(G_{a'/b'}) = [a' - 3 - (b' - 2)r, (b' - 1)r - a' + 3],$$

if  $r < p_l(a'/b')$  then

$$L_r(G_{a'/b'}) = \emptyset.$$

We construct the graph  $G_{p/q}$  as follows: Let  $X$  be the series parallel graph obtained from  $G_{a/b}$  and  $P_3$  by a parallel construction, and let  $Y$  be obtained from two copies of  $X$  by a series construction, let  $Z$  be obtained from  $Y$  and  $P_2$  by a parallel construction. Then  $G_{p/q}$  is obtained from  $G_{a'/b'}$  and  $Z$  by a series construction. The structure of  $G_{p/q}$  is as depicted in Fig. 4 below.




Figure 4: The graph  $G_{p/q}$

First we prove that if  $r < a/b$  then  $L_r(X) = \emptyset$  (and hence  $L_r(G) = \emptyset$ , as the sum or intersection of an emptyset with any other set is an emptyset). If  $r < p_l(a/b)$  then by the induction hypothesis,  $L_r(G_{a/b}) = \emptyset$ , hence  $L_r(X) = \emptyset$ . If  $p_l(a/b) \leq r < a/b$  then by the induction hypothesis,

$$L_r(G_{a/b}) = [a - 3 - (b - 2)r, (b - 1)r - a + 3].$$

So

$$L_r(X) = L_r(G_{a/b}) \cap L_r(P_3) = [a - 3 - (b - 2)r, (b - 1)r - a + 3] \cap [3 - r, 2r - 3].$$

Now it is straightforward to verify that  $L_r(X) = \emptyset$  (as  $r < a/b$ ).

Now we assume that  $a/b \leq r < \min\{8/3, (p - 2)/(q - 1)\}$ . By the induction hypothesis, for  $a/b \leq r < (a - 2)/(b - 1)$ ,

$$L_r(G_{a/b}) = [a - 3 - (b - 2)r, (b - 1)r - a + 3].$$

By Corollary 4.1,  $(p - 2)/(q - 1) \leq (a - 2)/(b - 1)$ , hence

$$\min\{8/3, (p - 2)/(q - 1)\} \leq \min\{8/3, (a - 2)/(b - 1)\}.$$

Therefore for  $a/b \leq r < \min\{8/3, (p-2)/(q-1)\}$ , we have

$$L_r(G_{a/b}) = [a-3-(b-2)r, (b-1)r-a+3]$$

and hence

$$L_r(X) = [3-r, (b-1)r-a+3] \cup [a-3-(b-2)r, 2r-3].$$

By Lemma 3.1

$$\begin{aligned} L_r(Y) &= L_r(X) + L_r(X) \\ &= [6-2r, (2b-2)r-2a+6] \cup [2a-6-2br+3r, 3r-6] \\ &\quad \cup [a-(b-1)r, br-a]. \end{aligned}$$

Since  $r < \min\{8/3, (a-2)/(b-1)\}$ , straightforward calculation shows that

$$L_r(Z) = L_r(Y) \cap L_r(P_2) = [a-(b-1)r, br-a].$$

Fig. 5 below indicates the positions of the intervals of  $L_r(Y)$  and  $L_r(P_2)$ .




Figure 5: The positions of  $L_r(Y)$  and  $L_r(P_2)$

It follows from the construction of  $G_{p/q}$  and Lemma 3.1 that  $L_r(G_{p/q}) = L_r(G_{a'/b'}) + L_r(Z)$ . Suppose  $s/t = p_l(a'/b')$ . By the induction hypothesis, for  $s/t \leq r < \min\{8/3, (a'-2)/(b'-1)\}$ ,

$$L_r(G_{a'/b'}) = [a'-3-(b'-2)r, (b'-1)r-a'+3].$$

It follows from the construction of the Farey sequence of order  $n$  that  $s/t \leq a/b$  (as  $s/t, a'/b'$  are two consecutive elements of row  $m$  for some  $m \leq n$ ). By Corollary 4.1,  $\min\{8/3, (p-2)/(q-1)\} \leq \min\{8/3, (a'-2)/(b'-1)\}$ . Therefore for  $a/b \leq r < \min\{8/3, (p-2)/(q-1)\}$ , we also have

$$L_r(G_{a'/b'}) = [a' - 3 - (b' - 2)r, (b' - 1)r - a' + 3].$$

Hence

$$\begin{aligned} L_r(G_{p/q}) &= L_r(Z) + L_r(G_{a'/b'}) \\ &= [a - (b - 1)r, br - a] + [a' - 3 - (b' - 2)r, (b' - 1)r - a' + 3] \\ &= [(a + a') - 3 - (b + b' - 2)r, (b + b' - 1)r - (a + a') + 3]. \end{aligned}$$

This completes the proof of Lemma 4.2. ■

**Proof of Theorem 2.1:** Suppose  $p/q \in (2, 8/3)$ . Let  $G_{p/q}$  be the graph constructed in the proof of Lemma 4.2. Let  $H_{p/q}$  be the graph obtained from  $G_{p/q}$  and  $P_3$  by a parallel construction. We shall show that  $\chi_c(H_{p/q}) = p/q$ .

If  $r < p/q$ , then either  $r < p_l(p/q)$  and hence  $L_r(G_{p/q}) = \emptyset$  which implies that  $L_r(H_{p/q}) = \emptyset$ , or  $p_l(p/q) \leq r < p/q$  and

$$L_r(G_{p/q}) = [p - 3 - (q - 2)r, (q - 1)r - p + 3].$$

In the latter case we have

$$\begin{aligned} L_r(H_{p/q}) &= L_r(G_{p/q}) \cap L_r(P_3) \\ &= [p - 3 - (q - 2)r, (q - 1)r - p + 3] \cap [3 - r, 2r - 3] \\ &= \emptyset. \end{aligned}$$

If  $r = p/q$ , then

$$\begin{aligned} L_r(H_{p/q}) &= L_r(G_{p/q}) \cap L_r(P_3) \\ &= [p - 3 - (q - 2)r, (q - 1)r - p + 3] \cap [3 - r, 2r - 3] \\ &= \{3 - r, 2r - 3\}. \end{aligned}$$

Therefore  $H_{p/q}$  is  $p/q$ -colourable, but not  $r$ -colourable for any  $r < p/q$ . So  $\chi_c(H_{p/q}) = p/q$ . This completes the proof of Theorem 2.1.

## References

- [1] J. A. Bondy and P. Hell, *A note on the star chromatic number*, J. Graph Theory **14** (1990), 479-482.

- [2] C. Chien and X. Zhu, *The circular chromatic number of series-parallel graphs with large girth*, Journal of Graph Theory, 33(2000), 185-198.
- [3] P. Hell and X. Zhu, *The circular chromatic number of series-parallel graphs*, Journal of Graph Theory, 33(2000), 14-24.
- [4] S. Liaw, Z. Pan and X. Zhu, *Construction of  $K_n$ -minor free graphs with given circular chromatic number*, Discrete Mathematics, to appear.
- [5] D. Moser, *The star chromatic number of planar graphs*, J. Graph Theory, 24(1997), 33-43.
- [6] I. Niven, H. Zuckerman and H. Montgomery, **An Introduction to the Theory of Numbers**, John Wiley & Sons, Inc., Fifth Edition, 1991.
- [7] J. G. Oxley, **Matroid Theory**, Oxford Science Publications, 1992.
- [8] Z. Pan and X. Zhu, *The circular chromatic number of series-parallel graph of large odd girth*, Discrete Mathematics, 245(2002), 235-246.
- [9] Z. Pan and X. Zhu, *Tight relation between the circular chromatic number and the girth of series-parallel graphs*, Discrete Mathematics, 254(2002), 393-404.
- [10] A. Vince, *Star chromatic number*, J. Graph Theory 12 (1988), 551-559.
- [11] X. Zhu, *Star chromatic numbers and products of graphs*, J. Graph Theory **16** (1992), 557-569.
- [12] X. Zhu, *Planar graphs with circular chromatic numbers between 3 and 4*, J. Combinatorial Theory Series B, **76** (1999), 170-200.
- [13] X. Zhu, *Circular chromatic number and graph minor*, Taiwanese Journal of Mathematics, 4(2000), 643-660.
- [14] X. Zhu, *Circular chromatic number, a survey*, Discrete Mathematics, Vol. 229 (1-3) (2001), 371-410.