

國立中山大學

NATIONAL SUN YAT-SEN UNIVERSITY

線性代數 (一)

MATH 103 / GEAI 1215: Linear Algebra I

期末考

January 6, 2020

Final Examination

姓名 Name : \_\_\_\_\_

學號 Student ID # : \_\_\_\_\_

Lecturer: Jephian Lin 林晉宏

Contents: cover page,  
**8 pages** of questions,  
score page at the end

To be answered: on the test paper

Duration: **110 minutes**

Total points: **30 points** + 7 extra points

**Do not open this packet until instructed to do so.**

Instructions:

- Enter your **Name** and **Student ID #** before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining it or circling it**. If multiple answers are shown then no marks will be awarded.
- 可用中文或英文作答

1. [1pt] Write down an example of a system of **linear** equations in variables  $a$ ,  $b$ , and  $c$ .
2. [1pt] Write down an example of a system of equations in variables  $a$ ,  $b$ , and  $c$  that is **not a linear system**.
3. [1pt] Write down an example of a system of **three linear equations** in its **echelon form** that contains **two free variables**.
4. [1pt] Write down an example of a  $4 \times 4$  **nonsingular** matrix.
5. [1pt] Write down an example of a  $4 \times 4$  **singular** matrix.

6. [1pt] Let  $S \subseteq \mathbb{R}^3$  be a set of vectors. Give an example of  $S$  such that  $S$  is linearly independent and  $\text{span}(S) \neq \mathbb{R}^3$ .
7. [1pt] Let  $S \subseteq \mathbb{R}^3$  be a set of vectors. Give an example of  $S$  such that  $\text{span}(S) = \mathbb{R}^3$  and  $S$  is not linearly independent.
8. [1pt] Let  $S \subseteq \mathbb{R}^3$  be a set of vectors. Give an example of  $S$  such that  $S$  is linearly independent and  $\text{span}(S) = \mathbb{R}^3$ .
9. [1pt] Let  $V \subseteq \mathbb{R}^3$  be a set of vectors. Give an example of  $V$  such that  $V$  is a subspace of  $\mathbb{R}^3$ .
10. [1pt] Let  $V \subseteq \mathbb{R}^3$  be a set of vectors. Give an example of  $V$  such that  $V$  is a not subspace of  $\mathbb{R}^3$ .

11. [1pt] Give an example of a function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $f$  is not a homomorphism.
  
  
  
  
  
  
  
  
  
  
12. [1pt] Give an example of a function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $f$  is a homomorphism but not an isomorphism.
  
  
  
  
  
  
  
  
  
  
13. [1pt] Give an example of a function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $f$  is an isomorphism.
  
  
  
  
  
  
  
  
  
  
14. [1pt] Suppose  $V_1$  and  $V_2$  are two subspaces of  $\mathbb{R}^3$ . Give an example of  $V_1$  and  $V_2$  such that they are not linearly independent (in terms of subspaces).
  
  
  
  
  
  
  
  
  
  
15. [1pt] Suppose  $V_1$  and  $V_2$  are two subspaces of  $\mathbb{R}^3$ . Give an example of  $V_1$  and  $V_2$  such that they are linearly independent (in terms of subspaces).

16. Let  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  and  $\mathcal{D} = \{\mathbf{u}_1, \mathbf{u}_2\}$  with

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{u}_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \text{ and } \mathbf{u}_2 = \begin{bmatrix} 3 \\ 2 \end{bmatrix}.$$

Define a homomorphism  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  such that  $f(\mathbf{v}_1) = 4\mathbf{u}_1$ ,  $f(\mathbf{v}_2) = 6\mathbf{u}_2$ , and  $f(\mathbf{v}_3) = 8\mathbf{u}_1 + 8\mathbf{u}_2$ .

(a) [2pt] Find  $\text{Rep}_{\mathcal{B}, \mathcal{D}}(f)$ .

(b) [3pt] Find a matrix  $A$  such that  $f(\mathbf{v}) = A\mathbf{v}$  for any  $\mathbf{v} \in \mathbb{R}^3$ .

17. [5pt] Let  $f : V \rightarrow W$  be a homomorphism. Show that  $f(X)$  is a subspace of  $W$  if  $X$  is a subspace of  $V$ .

18. [5pt] Let  $f : V \rightarrow W$  be a homomorphism. Show that  $f$  is one-to-one if and only if the null space of  $f$  is  $\{\mathbf{0}\}$ .

19. Let  $E_{ij}$  be the  $2 \times 3$  matrix whose entries are all zeros except that the  $i, j$ -entry is one. Then

$$\mathcal{B} = \{E_{11}, E_{12}, E_{13}, E_{21}, E_{22}, E_{23}\}$$

is a basis of  $\mathcal{M}_{2 \times 3}$ , the space of all  $2 \times 3$  real matrices. Suppose  $f : \mathcal{M}_{2 \times 3} \rightarrow \mathcal{M}_{2 \times 3}$  is a homomorphism such that  $\text{Rep}_{\mathcal{B}, \mathcal{B}}(f)$  equals

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

(a) [extra 1pt] Let  $M = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ . Find  $f(M)$ .

(b) [extra 2pt] Find the range of  $f$ .

(c) [extra 2pt] Find the nullspace of  $f$ .



20. [extra 2pt] Recall that  $\mathcal{L}(V, W)$  is the space of all homomorphisms from  $V$  to  $W$ . Let  $V = \mathcal{M}_{4 \times 5}$  be the space of all  $4 \times 5$  real matrices. Let  $W = \mathcal{P}_{100}$  be the space of all polynomials with real coefficients and of degree at most 100. Answer the following questions:

- (a) What is the zero vector in  $\mathcal{L}(V, W)$ ?
- (b) What is the dimension of  $V$ ?
- (c) What is the dimension of  $W$ ?
- (d) What is the dimension of  $\mathcal{L}(V, W)$ ?

**[END]**

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	5	
6	5	
7	5	
8	2	
Total	30 (+7)	