Bull. Aust. Math. Soc. **106** (2022), 288–300 doi:10.1017/S0004972721001076

# ON STARK'S CLASS NUMBER CONJECTURE AND THE GENERALISED BRAUER-SIEGEL CONJECTURE

## PENG-JIE WONG

(Received 13 October 2021; accepted 5 November 2021; first published online 10 January 2022)

#### Abstract

Stark conjectured that for any  $h \in \mathbb{N}$ , there are only finitely many CM-fields with class number h. Let C be the class of number fields L for which L has an almost normal subfield K such that L/K has solvable Galois closure. We prove Stark's conjecture for  $L \in C$  of degree greater than or equal to 6. Moreover, we show that the generalised Brauer–Siegel conjecture is true for asymptotically good towers of number fields  $L \in C$  and asymptotically bad families of  $L \in C$ .

2020 Mathematics subject classification: primary 11R29; secondary 11R42, 11R47.

*Keywords and phrases*: Stark's conjecture, Brauer–Siegel conjecture, class numbers, asymptotically exact families.

## 1. Introduction

For a number field  $L/\mathbb{Q}$ , let  $n_L$  and  $d_L$  denote its degree and absolute discriminant, respectively. Let  $h_L$  denote the class number of L. Stark [13] conjectured that for any  $h \in \mathbb{N}$ , there are only finitely many CM-fields, that is, totally imaginary quadratic extensions of totally real fields, with class number h. Stark proved his conjecture for fields with fixed degree greater than or equal to 6.

A number field *M* is called almost normal if there exists a sequence of number fields  $\mathbb{Q} = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_\ell = M$ , for some  $\ell \in \mathbb{N}$ , such that each extension  $M_i/M_{i-1}$  is normal for  $1 \le i \le \ell$ . In [11], Odlyzko verified Stark's conjecture for CM-fields *L* with  $n_L \ge 6$  whose maximal totally real subfields  $L^+$  are almost normal. In [6], Hoffstein and Jochnowitz proved that there exists an effective absolute constant *C* such that the assumption of the almost normality of  $L^+$  in Odlyzko's theorem can be replaced with the bound  $d_{L^+} > C^{[L^+:\mathbb{Q}]}$ . Murty [10] proved Stark's conjecture for CM-fields *L* having solvable normal closures over  $\mathbb{Q}$  when  $n_L \ge 6$ .

For any CM-field L, class field theory tells us that the class number  $h_{L^+}$  divides  $h_L$ , where  $L^+$  is the maximal totally real subfield of L. Thus, the number  $h_L^- = h_L/h_{L^+}$  is a

The author is currently an NCTS postdoctoral fellow.

<sup>©</sup> The Author(s), 2022. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

positive integer. This number is called the relative class number of *L*. One may also consider the following problem that generalises Stark's conjecture.

**PROBLEM 1.1.** Let *H* be a positive integer. Is the set of CM-fields *L* with  $h_L^- \leq H$  finite?

If the answer to this problem is affirmative, then Stark's conjecture is true. Since the class number of  $\mathbb{Q}$  is 1, Problem 1.1 for CM-fields *L* with  $n_L = 2$  is, in fact, Gauss's original class number conjecture proved by Deuring, Hecke, Heilbronn and Mordell. An effective version of Gauss's conjecture has been established by Goldfeld [4].

The first goal of this paper is to prove an instance of Problem 1.1. Let K be a subfield of a number field L. We say L/K has solvable Galois closure if  $Gal(\tilde{L}/K)$  is solvable, where  $\tilde{L}$  denotes the normal closure of L over K. Throughout this paper, we let C be the class of number fields L for which L has an almost normal subfield K such that L/K has solvable Galois closure. Our first main result is the following theorem.

## **THEOREM 1.2.** Let *H* be a positive integer.

(i) The set of CM-fields  $L \in C$  with  $n_L \neq 4$  and  $h_L^- \leq H$  is finite and its cardinality can be bounded effectively (in terms of H). Moreover, if Artin's conjecture is true, then the set of CM-fields L with  $n_L \neq 4$  and  $h_L^- \leq H$  is finite and its cardinality can be bounded effectively.

(ii) Unconditionally, the set of CM-fields L with  $n_L = 4$  and  $h_I^- \le H$  is finite.

**REMARK 1.3.** (i) The class *C* contains every number field with solvable Galois closure over  $\mathbb{Q}$ . If the maximal totally real subfield of *L* is almost normal, then *L* is also almost normal and so  $L \in C$ . Thus, the class *C* includes the fields considered by Murty [10] and Odlyzko [11] and Theorem 1.2 presents a common extension of their results. There is a further extension of Murty's result because the Galois closure of  $L \in C$  over  $\mathbb{Q}$  may not be solvable. For instance, if *L* contains a subfield *K* that is a nonsolvable Galois extension of  $\mathbb{Q}$ , then the Galois closure of *L* over  $\mathbb{Q}$  cannot be solvable even if the Galois closure of *L* over *K* is solvable.

(ii) The second part of Theorem 1.2 is ineffective since the proof requires Siegel's ineffective bound on the exceptional zeros of Dedekind zeta functions of quadratic fields.

There are several questions and results concerning the behaviour of  $h_L$  as L varies. Most famously, in [1], Brauer proved the following theorem.

**THEOREM 1.4 (Brauer–Siegel theorem)**. Let  $\{L_i\}$  be a family of number fields (that is,  $L_i \neq L_j$  for any  $i \neq j$ ) such that each  $L_i/\mathbb{Q}$  is Galois. If  $\lim_{i\to\infty} d_{L_i}^{1/n_{L_i}} = \infty$ , then

$$\lim_{i \to \infty} \frac{\log(h_{L_i} R_{L_i})}{\log \sqrt{d_{L_i}}} = 1$$

where  $R_{L_i}$  denotes the regulator of  $L_i$ . Moreover, the Galois condition on  $L_i/\mathbb{Q}$  can be removed if the generalised Riemann hypothesis is true.

Siegel [12] proved this theorem for the case when  $\{L_i\}$  is a family of quadratic fields. If  $\{L_i\}$  is a family of imaginary quadratic fields, then  $R_{L_i} = 1$  and hence the Brauer–Siegel theorem provides a rate at which  $h_{L_i}$  goes to infinity. Consequently, there are only finitely many imaginary quadratic fields with a bounded class number (see also [5]).

In [13], Stark proved an effective and much stronger version of the Brauer–Siegel theorem for families of almost normal number fields that do not contain any quadratic fields. Stark was aware that the Brauer–Siegel theorem would hold over families of almost normal fields, but this was only made explicit in the work of Zykin [15].

In [14], Tsfasman and Vlăduţ formulated the generalised Brauer–Siegel conjecture (GBS) and proved some instances of it. Let  $\mathcal{L} = \{L_i\}$  be a family of number fields. By the definition of the Dedekind zeta function  $\zeta_{L_i}(s)$  of  $L_i$ , there are nonnegative integers  $N_a(L_i)$  such that

$$\zeta_{L_i}(s) = \prod_q (1 - q^{-s})^{-N_q(L_i)},$$

for Re(s) > 1, where the product is over prime powers *q*. We call  $\mathcal{L} = \{L_i\}$  asymptotically exact if the limits

$$\phi_q(\mathcal{L}) = \lim_{i \to \infty} \frac{N_q(L_i)}{g_{L_i}}, \quad \phi_{\mathbb{R}}(\mathcal{L}) = \lim_{i \to \infty} \frac{r_1(L_i)}{g_{L_i}} \quad \text{and} \quad \phi_{\mathbb{C}}(\mathcal{L}) = \lim_{i \to \infty} \frac{r_2(L_i)}{g_{L_i}}$$

exist, for every prime power q, where  $g_{L_i} = \log \sqrt{d_{L_i}}$ , and  $r_1(L_i)$  and  $r_2(L_i)$  are the numbers of real and complex embeddings of  $L_i$ , respectively.

CONJECTURE 1.5 (Generalised Brauer–Siegel conjecture). For any asymptotically exact family  $\mathcal{L} = \{L_i\}$ , the limit  $BS(\mathcal{L}) = \lim_{i \to \infty} \log(h_{L_i}R_{L_i})/g_{L_i}$  exists. Moreover,

$$BS(\mathcal{L}) = 1 + \sum_{q} \phi_q(\mathcal{L}) \log \frac{q}{q-1} - \phi_{\mathbb{R}}(\mathcal{L}) \log 2 - \phi_{\mathbb{C}}(\mathcal{L}) \log 2\pi.$$

Following [14], an asymptotically exact family  $\mathcal{L}$  will be called asymptotically bad if  $\phi_q(\mathcal{L}) = \phi_{\mathbb{R}}(\mathcal{L}) = \phi_{\mathbb{C}}(\mathcal{L}) = 0$  for every prime power q. If an asymptotically exact family is not asymptotically bad, then it will be said to be asymptotically good.

Tsfasman and Vlăduţ [14] proved their conjecture for any asymptotically exact family under the assumption of the generalised Riemann hypothesis. Also, unconditionally, they proved GBS for asymptotically good towers of almost normal fields. (Recall that a family { $L_i$ } is called a tower if  $L_i \subsetneq L_{i+1}$  for each *i*.) Moreover, Zykin [15] proved GBS for asymptotically bad families of almost normal fields. Recently, Dixit [3] showed that GBS is true for asymptotically good towers and asymptotically bad families of number fields with solvable Galois closure over  $\mathbb{Q}$ . Our second goal is to prove the following extension of the work of Dixit, Tsfasman and Vlăduţ, and Zykin.

THEOREM 1.6. If an exact family  $\mathcal{L} \subseteq C$  is an asymptotically good tower or an asymptotically bad family, then GBS is true for  $\mathcal{L}$ . Furthermore, if Artin's conjecture

is true, then GBS is true for any asymptotically good tower and any asymptotically bad family.

**REMARK** 1.7. (i) As noted by Dixit [3], it follows from the class number formula that GBS for an exact family  $\mathcal{L} = \{L_i\}$  is equivalent to the statement that the limit

$$\rho(\mathcal{L}) = \lim_{i \to \infty} \frac{\log \rho_{L_i}}{g_{L_i}}$$

exists and equals  $\sum_{q} \phi_q(\mathcal{L}) \log(q/(q-1))$ , where  $\rho_{L_i}$  is the residue of the Dedekind zeta function  $\zeta_{L_i}(s)$  at s = 1. We shall use this fact repeatedly in Section 4.

(ii) The family  $\mathcal{L} = \{L_i\}$  is asymptotically bad if and only if  $\lim_{i\to\infty} d_{L_i}^{1/n_{L_i}} = \infty$ . From this, GBS for asymptotically bad families is equivalent to the classical Brauer–Siegel conjecture (that is, the Brauer–Siegel 'theorem' without the Galois condition on  $L_i$ ).

## 2. Stark zeros in certain towers of number fields

Stark's conjecture arose from his work on exceptional zeros of Dedekind zeta functions. These zeros are also crucial to the study of GBS. In this section, we will briefly review some results concerning 'Stark zeros' (see [10] for further discussion).

For any number field *L*, Stark showed that the Dedekind zeta function  $\zeta_L(s)$  has at most one zero in the region

$$1 - \frac{1}{4\log d_L} \le \operatorname{Re}(s) \le 1$$
 and  $|\operatorname{Re}(s)| < \frac{1}{4\log d_L}$ , (2.1)

and if the zero exists in this region, it must be real and simple. We call this possible zero the Stark zero of  $\zeta_L(s)$  and denote it by  $\beta_0$ . For any Galois extension L/K, if the Stark zero  $\beta_0$  of  $\zeta_L(s)$  exists, then there is a field M with  $K \subseteq M \subseteq L$  such that  $[M : K] \leq 2$  and  $\zeta_M(\beta_0) = 0$ . In the case when L is almost normal, Stark [13, Lemma 10] showed that if the Stark zero  $\beta_0$  of  $\zeta_L(s)$  appears in the region

$$1 - \frac{1}{16 \log d_L} \le \operatorname{Re}(s) \le 1,$$

then there is a quadratic field  $F \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ . These results play a key role in the argument of Odlyzko [11].

Let *m* be a positive integer. We set

$$e(m) = \max_{p^{\alpha} \parallel m} \alpha$$
 and  $\delta(m) = (e(m) + 1)^2 3^{1/3} 12^{e(m)-1}$ . (2.2)

THEOREM 2.1 (Murty, [10, Theorem 2.1]). Let L/K be an extension with solvable Galois closure. Let m be the degree of L/K. There exists an absolute constant  $c_0 > 0$  such that if  $\zeta_L(s)$  has a zero  $\beta_0$  in the region

$$1 - \frac{c_0}{m^{e(m)}\delta(m)\log d_L} \le \operatorname{Re}(s) \le 1 \quad and \quad |\operatorname{Re}(s)| < \frac{c_0}{m^{e(m)}\delta(m)\log d_L},$$

then there is a field M with  $K \subseteq M \subseteq L$  such that  $[M : K] \leq 2$  and  $\zeta_M(\beta_0) = 0$ .

As in the work of Odlyzko [11], in order to 'descend' the Stark zero of  $\zeta_L(s)$  to a quadratic field, Murty considered the class of number fields with solvable Galois closure over  $\mathbb{Q}$  and applied his theorem with  $K = \mathbb{Q}$ .

We require the following refinement of Murty's theorem for our main theorems.

**THEOREM 2.2.** Let L/K be an extension with solvable Galois closure and let K be an almost normal field. Let m and n be the degrees of L/K and  $L/\mathbb{Q}$ , respectively. Let  $c_0$  be as in Theorem 2.1 and  $c_1 = \min\{c_0, 1/16\}$ . Suppose that  $\zeta_L(s)$  has a zero  $\beta_0$  in the region

$$1 - \frac{c_1}{m^{e(m)}\delta(m)\log d_L} \le \operatorname{Re}(s) \le 1 \quad and \quad |\operatorname{Re}(s)| < \frac{1}{4\log d_L}, \tag{2.3}$$

where e(m) and  $\delta(m)$  are defined as in (2.2). Then  $\beta_0$  is real and simple and there is a quadratic field  $F \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ . Consequently, if  $\zeta_L(s)$  has a zero  $\beta_0$  in the region

$$1 - \frac{c_1}{n^{e(n)}\delta(n)\log d_L} \le \operatorname{Re}(s) \le 1 \quad and \quad |\operatorname{Re}(s)| < \frac{1}{4\log d_L},\tag{2.4}$$

then  $\beta_0$  is real and simple and there is a quadratic field  $F \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ .

**PROOF.** As the region (2.3) is contained in the region (2.1),  $\beta_0$  must be real and simple. Since  $c_1 \le c_0$ , by Theorem 2.1, there is a field M with  $K \subseteq M \subseteq L$  such that  $[M : K] \le 2$  and  $\zeta_M(\beta_0) = 0$ . As  $[M : K] \le 2$ , the extension M/K is normal. Since K is almost normal, M is also almost normal. Observing that

$$\frac{c_1}{m^{e(m)}\delta(m)\log d_L} \le \frac{1}{16\log d_M},$$

we have  $\beta_0 \in [1 - 1/16 \log d_M, 1]$ . Finally, by the result on Stark zeros in almost normal extensions [13, Lemma 10], there is a quadratic field *F* with  $F \subseteq M \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ .

To prove the second part of the theorem, it suffices to show that the region (2.4) is contained in the region (2.3). As  $m \mid n$ , we have  $e(m) \leq e(n)$  and thus  $\delta(m) \leq \delta(n)$ . This concludes the proof.

If Artin's conjecture is true, one has the following improvement of these results.

LEMMA 2.3 (Stark, [13, Lemma 13]). Let *L* be a number field of degree n > 1 and let  $\tilde{L}$  be the normal closure of *L* over  $\mathbb{Q}$ . Suppose that all the Artin *L*-functions  $L(s, \psi, \tilde{L}/\mathbb{Q})$  are holomorphic at  $s \neq 1$ . If there is a zero  $\beta_0$  of  $\zeta_L(s)$  in the range

$$1 - \frac{1}{8(n-1)\log d_L} \le \operatorname{Re}(s) \le 1,$$

then there exists a quadratic field  $F \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ .

Lastly, we recall the following bound for the exceptional zeros of Dedekind zeta functions of quadratic fields.

LEMMA 2.4 [13, Lemma 11]. Let *F* be a quadratic field. Then there is an effective absolute constant  $c_2 > 0$  such that  $\zeta_F(\sigma)$  is nonvanishing for  $\sigma \ge 1 - c_2/d_F^{1/2}$ .

#### 3. Proof of Theorem 1.2

To prove Theorem 1.2, we shall prove the following lower bounds for  $h_{I}^{-}$ .

THEOREM 3.1. Let L be a CM-field with maximal totally real subfield  $L^+$ . Assume either  $L \in C$  or the truth of Artin's conjecture. Write  $d_L = d_{L^+}^2 f$  and set  $n = [L^+ : \mathbb{Q}]$ . Then there are effective absolute constants  $c_3 > 0$  and  $c_4 > 1$  such that

$$h_L^- \ge \frac{c_3}{ng(n)} c_4^n f^{(1/2) - (1/2n)},$$
(3.1)

where the function g(n) is defined by

$$g(n) = \begin{cases} (2n)^{e(2n)}\delta(2n) & \text{if } L \in C, \\ n - \frac{1}{2} & \text{if } Artin's \ conjecture \ is \ true. \end{cases}$$

Moreover, for any  $\varepsilon \in [1/(8 \log d_L), 1]$ , there exists  $c(\varepsilon) > 0$ , depending only on  $\varepsilon$ , such that

$$h_{L}^{-} \ge \frac{c_{3}c(\varepsilon)^{n}}{ng(n)} d_{L^{+}}^{(1/2) - (1/n) - \varepsilon} f^{(1/2) - (1/2n)}.$$
(3.2)

**PROOF.** We shall follow the arguments used in [10, 11, 13]. Note that there is a nontrivial Hecke character  $\chi$  of  $L^+$  such that  $\zeta_L(s) = \zeta_{L^+}(s)L(s,\chi)$ , where  $L(s,\chi)$  is the Hecke *L*-function attached to  $\chi$ . Let  $\beta_0$  be the possible exceptional zero of  $L(s,\chi)$  satisfying

$$1 - \frac{1}{4\log(d_{I^+}^2 f)} \le \beta_0 \le 1.$$

and set  $\beta_1 = \beta_0$  if it exists. Otherwise, set  $\beta_1 = 1 - 1/(4 \log(d_{L^+}^2 f))$ . By equation (31) in [13],

$$h_L^- = \frac{h_L}{h_{L^+}} \ge d_{L^+}^{1/2} f^{1/2} (2\pi)^{-n} L(1,\chi).$$

In addition, as in [13, Proof of Lemma 5] (see also [11, Equation (4.2)]), for any  $\sigma_1$  such that

$$1 + \frac{1}{4\log d_L} \le \sigma_1 \le 2,$$

there is a  $c_5 > 0$  such that

$$L(1,\chi) \ge c_5(1-\beta_1)d_{L^+}^{-(1/2)(\sigma_1-1)}\zeta_{L^+}(\sigma_1)^{-1}.$$

Thus,

$$h_{L}^{-} \ge c_{5}(1-\beta_{1})d_{L^{+}}^{(1/2)-(1/2)(\sigma_{1}-1)}f^{1/2}(2\pi)^{-n}\zeta_{L^{+}}(\sigma_{1})^{-1}.$$
(3.3)

Now, if  $L(s, \chi)$  has a zero  $\beta_0$  in the region

$$1 - \frac{c_1}{g(n)\log d_L} \le \operatorname{Re}(s) \le 1, \tag{3.4}$$

[7]

where  $c_1$  is defined as in Theorem 2.2, then  $\beta_0$  is also a zero of  $\zeta_L(s)$ . By Theorem 2.2 and Lemma 2.3, since  $\beta_0$  is in the region (3.4) and  $c_1 \leq 1/16$ , there is a quadratic field  $F \subseteq L$  such that  $\zeta_F(\beta_0) = 0$ . By Lemma 2.4,  $\beta_0 \leq 1 - c_2/d_F^{1/2}$ . Therefore,  $L(s, \chi)$ is nonzero on

$$\max\left\{1 - \frac{c_1}{g(n)\log d_L}, 1 - \frac{c_2}{d_L^{1/2n}}\right\} \le \operatorname{Re}(s) \le 1.$$

(Here, we used  $d_L \ge d_{F^*}^n$ ) Consequently, as  $d_L = d_{L^*}^2 f$ ,

$$1 - \beta_1 \ge \frac{c_6}{g(n)} \min\{(\log d_L)^{-1}, d_L^{-1/2n}\} \ge \frac{c_6}{ng(n)} d_{L^+}^{-1/2n} f^{-1/2n},$$

where  $c_6 = \min\{c_1, c_2\} > 0$ . It follows from (3.3) that

$$h_{L}^{-} \geq \frac{c_{5}c_{6}}{ng(n)} d_{L^{+}}^{(1/2)-(1/2)(\sigma_{1}-1)-(1/n)} f^{(1/2)-(1/2n)} (2\pi)^{-n} \zeta_{L^{+}} (\sigma_{1})^{-1}.$$
(3.5)

Now, as argued by Odlyzko [11, pages 284–285], there are effective absolute positive constants  $c_7$  and  $c_8$  such that

$$h_L^- \ge \frac{c_5 c_6 c_7}{ng(n)} f^{(1/2) - (1/2n)} (1 + c_8)^n.$$

This proves the first claimed bound.

To prove the second bound, we choose  $\sigma_1 = 1 + 2\varepsilon$  with  $\varepsilon \in [1/(8 \log d_L), 1]$ . Recall that for  $\sigma \in (1, 3]$ ,

$$0 < \zeta_{L^+}(\sigma) \le \zeta(\sigma)^n \le \left(\frac{c_9}{\sigma - 1}\right)^n$$

for some effective absolute  $c_9 > 0$ . Thus, by (3.5),

$$h_{L}^{-} \geq \frac{c_{5}c_{6}}{ng(n)} d_{L^{+}}^{(1/2)-(1/n)-\varepsilon} f^{(1/2)-(1/2n)} (2\pi)^{-n} \zeta (1+2\varepsilon)^{-n}$$
  
$$\geq \frac{c_{5}c_{6}}{ng(n)} d_{L^{+}}^{(1/2)-(1/n)-\varepsilon} f^{(1/2)-(1/2n)} \left(\frac{\varepsilon}{c_{9}\pi}\right)^{n}$$

as desired.

We are now in a position to prove Theorem 1.2. As mentioned before, for CM-fields of degree 2, the theorem follows from the works of Goldfeld, Gross and Zagier. So, we may restrict our attention to CM-fields of degree at least 6.

**PROOF OF THEOREM 1.2.** (i) Let *L* be a CM-field with  $h_L^- \le H$ ,  $n_L + \ge 3$  and  $d_L = d_{L^+}^2 f$ . Assume either  $L \in C$  or the truth of Artin's conjecture. Since  $e(n) \le \log n/\log 2$ ,

$$n^{e(n)} \le \exp((\log n)^2).$$

294

Thus, by (3.1), there exists an effective positive constant  $A_H > 0$ , depending only on H, such that  $n_{L^+} \le A_H$  and  $f \le A_H$ .

Note that  $n_L \ge 6$  for  $[L^+ : \mathbb{Q}] > 2$ . By Minkowski's bound,  $n_L \le 2 \log d_L / \log 3$ , and thus  $1/(8 \log d_L) \le 1/(4(\log 3)n_L) \le 1/(24 \log 3)$ . To use (3.2), we may choose  $\varepsilon = 1/(24 \log 3)$  so that

$$\frac{1}{2} - \frac{1}{n_{L^+}} - \varepsilon \ge \frac{1}{6} - \varepsilon > 0.$$

Now, by (3.2) and the bounds  $n_{L^+} \le A_H$  and  $f \le A_H$ , there is an effective positive constant  $B_H > 0$ , depending only on H, such that  $d_{L^+} \le B_H$ . Hence, we arrive at  $d_L = d_{L^+}^2 f \le B_H^2 A_H$ . This completes the proof because the number of number fields L with  $d_L \le B_H^2 A_H$  can be bounded effectively.

(ii) In the proof of Theorem 3.1, instead of using Lemma 2.4, we may use Siegel's ineffective bound

$$\beta_0 \le 1 - \frac{\tilde{c}_2}{d_F^{1/4}}$$

for some ineffective absolute  $\tilde{c}_2 > 0$ . This results in an ineffective improvement of the second bound of Theorem 3.1 to give

$$h_L^- \ge \frac{\tilde{c}_{3C}(\varepsilon)^n}{ng(n)} d_{L^+}^{(1/2) - (1/2n) - \varepsilon} f^{(1/2) - (1/4n)}$$

for some ineffective  $\tilde{c}_3 > 0$ . This estimate, together with the first bound of Theorem 3.1, proves the finiteness of *L* with  $n_{L^+} = 2$  and  $h_L^- \le H$  for any given  $H \in \mathbb{N}$ . Note that *L* must have Galois closure over  $\mathbb{Q}$  since  $n_L = 4$  and so this finiteness result is unconditional.

## 4. Proof of Theorem 1.6

**4.1. Auxiliary lemmas.** Our proof of Theorem 1.6 builds on the works of Dixit [3], Tsfasman and Vlăduț [14] and Zykin [15].

Let *L* be a number field and let  $G_L(s)$  be the entire function such that

$$\zeta_L(s) = \frac{\rho_L}{s-1} G_L(s). \tag{4.1}$$

We shall set

$$Z_L(s) = \frac{d}{ds} \left( \frac{\log G_L(s)}{g_L} \right)$$

From Lagarias and Odlyzko [8, Theorem 9.2], based on an idea of Tsfasman and Vlăduţ [14], Dixit proved the following lemma.

LEMMA 4.1 (Dixit, [3, Lemma 4.2]). Let *L* belong to an asymptotically good family  $\mathcal{L}$ , and let  $n = [L : \mathbb{Q}]$ . Suppose that  $\zeta_L(s)$  has no zero in the region

$$1 - \frac{c_1}{n^{e(n)}\delta(n)\log d_L} \le \operatorname{Re}(s) \le 1.$$
(4.2)

Then there are positive constants  $C_1$ ,  $C_2$  and  $C_3$ , dependent on  $c_1$  and  $\mathcal{L}$ , but independent of L, such that  $|Z_L(1 + \theta)| \leq C_1 g_L^{C_2 \log g_L}$  for any  $\theta \in (0, 1)$  and any  $g_L > C_3$ .

To control the possible exceptional zeros arising from an asymptotically good family, Tsfasman and Vlăduţ [14] proved the following finiteness lemma.

LEMMA 4.2 (Tsfasman and Vlăduţ, [14, Lemma 7.3]; see also [3, Lemma 4.3]). Let  $\mathcal{L} = \{L_i\}$  be an asymptotically good family, and set

$$Q(\mathcal{L}) = \{F \mid [F : \mathbb{Q}] = 2 \text{ and } F \subseteq L_i \text{ for some } i\}.$$

Then  $Q(\mathcal{L})$  is finite.

Lastly, we require the following bounds for the residues  $\rho_L$ .

LEMMA 4.3 (Louboutin, [9, Theorem 1]). For any number field  $L \neq \mathbb{Q}$ , if  $\zeta_L(\beta) = 0$  for some  $\beta \in [\frac{1}{2}, 1)$ , then

$$\rho_L \le (1-\beta) \Big( \frac{e \log d_L}{2n_L} \Big)^{n_L}.$$

LEMMA 4.4 (Stark, [13, Lemma 4]). There exists an effective constant c' > 0 such that for any number field  $L \neq \mathbb{Q}$ , one has  $\rho_L > c'(1 - \beta_0)$ , where  $\beta_0$  is the possible exceptional zero of  $\zeta_L(s)$  in  $[1 - 1/(4 \log d_L), 1]$ . If the possible exceptional zero does not exist, then one can take  $\beta_0 = 1 - 1/(4 \log d_L)$ .

**4.2. Proof of Theorem 1.6 when**  $\mathcal{L}$  **is an asymptotically good tower.** In light of the work of Dixit [3] and Tsfasman and Vlăduţ [14], to prove Theorem 1.6, we require the following general inequalities towards GBS for asymptotically good towers.

LEMMA 4.5. Let  $\mathcal{L} = \{L_i\}$  be an asymptotically good tower. Let  $\{\theta_{L_i}\} \subset (0, 1)$  be a sequence convergent to zero such that  $\log \theta_{L_i} = o(g_{L_i})$ . Then

$$\sum_{q} \phi_q(\mathcal{L}) \log \frac{q}{q-1} \ge \limsup_{i \to \infty} \frac{\log \rho_{L_i}}{g_{L_i}}$$

and

$$\liminf_{i\to\infty}\frac{\log\rho_{L_i}}{g_{L_i}}\geq \sum_q\phi_q(\mathcal{L})\log\frac{q}{q-1}-\limsup_{i\to\infty}\frac{\log G_{L_i}(1+\theta_{L_i})}{g_{L_i}}.$$

**PROOF.** The first inequality is established in [14] for any asymptotically exact family, so it remains to prove the second inequality. By (4.1), for any  $\theta_L \in (0, 1)$ ,

$$\frac{\log \zeta_L(1+\theta_L)}{g_L} = \frac{\log \rho_L}{g_L} + \frac{\log G_L(1+\theta_L)}{g_L} - \frac{\log \theta_L}{g_L}.$$
(4.3)

Since  $\mathcal{L} = \{L_i\}$  is a tower, for  $\theta > 0$ ,

Stark's class number conjecture

$$\begin{aligned} \frac{\log \zeta_{L_i}(1+\theta)}{g_{L_i}} &= \sum_q \frac{N_q(L_i)}{g_{L_i}} \log \frac{1}{1-q^{-1-\theta}} \\ &= \sum_p \frac{N_p(L_i)}{g_{L_i}} \log \frac{1}{1-p^{-1-\theta}} + \sum_{\substack{q=p^k \\ k \ge 2}} \frac{N_q(L_i)}{g_{L_i}} \log \frac{1}{1-q^{-1-\theta}} \\ &\ge \sum_p \phi_p(\mathcal{L}) \log \frac{1}{1-p^{-1-\theta}} + \sum_{\substack{q=p^k \\ k \ge 2}} \frac{N_q(L_i)}{g_{L_i}} \log \frac{1}{1-q^{-1-\theta}}.\end{aligned}$$

In addition, there is a positive number  $\delta$  such that

$$\sum_{\substack{q=p^k\\k\geq 2}} \frac{N_q(L_i)}{g_{L_i}} \log \frac{1}{1-q^{-1-\theta}} \to \sum_{\substack{q=p^k\\k\geq 2}} \phi_q(\mathcal{L}) \log \frac{1}{1-q^{-1-\theta}}$$

uniformly in  $\theta > -\delta$ . Thus, as the sequence  $\{\theta_{L_i}\}$  converges to zero, we arrive at

$$\liminf_{i \to \infty} \frac{\log \zeta_{L_i}(1+\theta_{L_i})}{g_{L_i}} \ge \sum_q \phi_q(\mathcal{L}) \log \frac{q}{q-1}.$$
(4.4)

Gathering (4.3), (4.4) and the assumption  $\log \theta_{L_i} = o(g_{L_i})$  completes the proof.

We now prove Theorem 1.6 for asymptotically good towers.

**PROOF OF THEOREM 1.6 FOR ASYMPTOTICALLY GOOD TOWERS.** Let  $\mathcal{L} = \{L_i\}$  be an asymptotically good tower. Assume either  $\mathcal{L} = \{L_i\} \subseteq C$  or Artin's conjecture. Following Dixit [3, Section 5.1], we choose  $\theta_{L_i} = g_{L_i}^{-(C_2+1)\log g_{L_i}}$ , where  $C_2$  is the same as in Lemma 4.1. It is clear that the sequence  $\{\theta_{L_i}\}$  converges to zero and  $\log \theta_{L_i} = o(g_{L_i})$ . Therefore, by Lemma 4.5, it remains to show that

$$\limsup_{i\to\infty}\frac{\log G_{L_i}(1+\theta_{L_i})}{g_{L_i}}\leq 0.$$

By Theorem 2.2 and Lemma 2.3, if  $\zeta_{L_i}(s)$  has a zero  $\beta_0$  in the region (4.2) with  $L = L_i$  and  $n = n_{L_i}$ , then it must be a real zero of  $\zeta_F(s)$  for some  $F \in Q(\mathcal{L})$ . By Lemmas 2.4 and 4.2,

$$\beta_0 \leq 1 - \frac{c_2}{\max_{F \in \mathcal{Q}(\mathcal{L})} d_F^{1/2}} < 1.$$

Hence, for *i* sufficiently large,  $\zeta_{L_i}(s)$  has no zero in the region (4.2) with  $L = L_i$  and  $n = n_{L_i}$ . As  $G_{L_i}(1) = 1$ , by Lemma 4.1, for *i* sufficiently large,

$$\frac{\log G_{L_i}(1+\theta_{L_i})}{g_{L_i}} = \int_0^{\theta_{L_i}} Z_{L_i}(1+\theta) \, d\theta \ll g_{L_i}^{-\log g_{L_i}}$$

Taking the lim sup on both sides completes the proof.

[10]

**4.3. Proof of Theorem 1.6 when**  $\mathcal{L}$  is an asymptotically bad family. For any asymptotically bad family  $\mathcal{L} = \{L_i\}$ , we have  $0 \ge \limsup_{i \to \infty} (\log \rho_{L_i})/g_{L_i}$ . Thus, to prove GBS for an asymptotically bad family  $\mathcal{L}$ , it is sufficient to show that

$$\liminf_{i \to \infty} \frac{\log \rho_{L_i}}{g_{L_i}} \ge 0 \tag{4.5}$$

Let I be the set of indices *i* for which  $\zeta_{L_i}(s)$  does not have any zero in the region (4.2) with  $L = L_i$  and  $n = n_{L_i}$ , and let  $\mathcal{J}$  be the set of indices *i* for which  $\zeta_{L_i}(s)$  has a zero in the region (4.2) with  $L = L_i$  and  $n = n_{L_i}$ . Without loss of generality, we may assume that both I and  $\mathcal{J}$  are infinite.

If  $\zeta_{L_i}(s)$  is nonvanishing on the region (4.2), Lemma 4.4 gives

$$\rho_{L_i} > c' \min\left\{\frac{c_1}{n_{L_i}^{e(n_{L_i})}\delta(n_{L_i})\log d_{L_i}}, \frac{1}{4\log d_{L_i}}\right\} \ge \frac{c'c_1}{n_{L_i}^{e(n_{L_i})}\delta(n_{L_i})\log d_{L_i}}$$

and thus we obtain

$$\frac{\log \rho_{L_i}}{g_{L_i}} \geq \frac{\log(c'c_1) - e(n_{L_i})\log n_{L_i} - \log \delta(n_{L_i}) - \log \log d_{L_i}}{g_{L_i}}$$

Recall the estimates  $e(n) \le \log n / \log 2$  and  $\delta(n) \ll n^4$ . Since  $g_{L_i} \to \infty$  and  $n_{L_i}/g_{L_i} \to 0$ , by the above inequality, we obtain

$$\liminf_{\substack{i\to\infty\\i\in I}}\frac{\log\rho_{L_i}}{g_{L_i}}\geq 0$$

Now suppose that  $\zeta_{L_i}(s)$  has a zero  $\beta_{0,i}$  in the region (4.2) with  $L = L_i$  and  $n = n_{L_i}$ . By Theorem 2.2 and Lemma 2.3, there is a quadratic field  $F_i \subseteq L_i$  such that  $\zeta_{F_i}(\beta_{0,i}) = 0$ . By Lemmas 4.3 and 4.4,

$$\rho_{L_i} = \rho_{L_i} \frac{\rho_{F_i}}{\rho_{F_i}} \ge c'(1 - \beta_{0,i}) \frac{1}{(1 - \beta_{0,i})(e \log d_{F_i}/2n_{F_i})^{n_{F_i}}} \rho_{F_i}$$

Hence,  $(\log \rho_{L_i})/g_{L_i}$  is bounded below by

$$\frac{\log c' + 2\log(4/e\log d_{F_i}) + \log \rho_{F_i}}{g_{L_i}} = \frac{\log c' + 2\log(4/e\log d_{F_i})}{g_{L_i}} + \frac{g_{F_i}}{g_{L_i}}\frac{\log \rho_{F_i}}{g_{F_i}}$$

By the classical Brauer–Siegel theorem, Theorem 1.4, for quadratic fields  $F_i$ ,

$$\liminf_{\substack{i \to \infty \\ i \in \mathcal{T}}} \frac{\log \rho_{L_i}}{g_{L_i}} \ge 0$$

Since  $I \cup \mathcal{J}$  contains all the indices, this gives (4.5) and completes the proof.

## 5. Concluding remarks

In [11], Odlyzko also showed that the generalised Riemann hypothesis implies Stark's conjecture. We note that the generalised Riemann hypothesis can be replaced by a much weaker conjecture on the nonexistence of Stark zeros. More precisely, assume that none of the Dedekind zeta functions  $\zeta_L(s)$  of CM-fields *L* admits a zero in the region (2.1). Then arguing as in the proof of Theorem 3.1, for any CM-field *L* with  $n_{L^+} = n$  and applying Minkowski's bound,

$$h_{L}^{-} \geq \frac{c_{5}c_{7}}{4} \frac{1}{\log d_{L}} d_{L^{+}}^{1/n} f^{1/2} c_{4}^{n} \geq \frac{c_{5}c_{7}\sqrt{3}}{4} \frac{1}{\log(d_{L^{+}}^{2}f)} f^{1/2} c_{4}^{n}$$

and

$$h_L^- \ge \frac{c_5 c(\varepsilon)^n}{4} \frac{1}{\log(d_{L^+}^2 f)} d_{L^+}^{(1/2)-\varepsilon} f^{1/2}$$

Consequently, the cardinality of the set of CM-fields L with  $h_L^- \le H$  is finite and can be bounded effectively (in terms of H).

In a slightly different vein, for GBS, suppose that a family  $\mathcal{L} = \{L_i\}$  is an asymptotically good tower or an asymptotically bad family such that each Dedekind zeta function  $\zeta_{L_i}(s)$  is nonzero in the region (2.1) with  $L = L_i$  and  $n = n_L$ . By the arguments used in Section 4 and observing that the region (4.2) is contained in the region (2.1), one can show that GBS is true for  $\mathcal{L}$ .

Last but not least, we note that Theorem 2.2 may have other applications. For instance, consider the Euler–Kronecker constant  $\gamma_L$  attached to a number field *L* in the expansion

$$\frac{\zeta'_L}{\zeta_L}(s) = \frac{-1}{s-1} + \gamma_L + O(s-1).$$

Ihara [7] showed that under the generalised Riemann hypothesis,  $\gamma_L \leq 2 \log \log \sqrt{d_L}$ , and unconditionally,  $\gamma_L \geq -\log \sqrt{d_L}$ . In [2], Dixit proved that for every almost normal *L* not containing any quadratic fields,  $|\gamma_L| \leq \kappa_0 (\log d_L)^4 n_L^3$  for some absolute  $\kappa_0 > 0$ . Moreover, he showed that if *L* has solvable Galois closure over  $\mathbb{Q}$  and does not contain any quadratic fields, then  $|\gamma_L| \leq \kappa_1 (\log d_L)^{\kappa_2 \log \log d_L}$  for some absolute  $\kappa_1 > 0$ and  $\kappa_2 > 0$ . Now, if we consider a number field  $L \in C$  of degree *n* such that *L* does not contain any quadratic fields, then by Theorem 2.2, any zero  $\beta_0$  of  $\zeta_L(s)$  in the region (2.1) must satisfy

$$1 - \frac{1}{4 \log d_L} \le \beta_0 \le 1 - \frac{c_1}{n^{e(n)} \delta(n) \log d_L}$$

Hence, arguing as in [2, Sections 2.3.1–2.3.2] yields the following theorem.

THEOREM 5.1. Let  $L \in C$  be a number field not containing any quadratic fields. Then

$$|\gamma_L| \le \kappa_1' (\log d_L)^{\kappa_2' \log \log d_L}$$

for some absolute  $\kappa'_1 > 0$  and  $\kappa'_2 > 0$ .

## Acknowledgement

The author would like to thank the referee for helpful comments.

### References

- [1] R. Brauer, 'On zeta-functions of algebraic number fields', Amer. J. Math. 2 (1947), 243–250.
- [2] A. B. Dixit, 'On Euler–Kronecker constants and the generalized Brauer–Siegel theorem', *Proc. Amer. Math. Soc.* **148** (2020), 1399–1414.
- [3] A. B. Dixit, 'On the generalized Brauer–Siegel theorem for asymptotically exact families with solvable Galois closure', *Int. Math. Res. Not. IMRN* 2021 (2021), 10941–10956.
- [4] D. Goldfeld, 'Gauss' class number problem for imaginary quadratic fields', *Bull. Amer. Math. Soc.* (*N.S.*) 13 (1985), 23–37.
- [5] H. A. Heilbronn, 'On the class number of imaginary quadratic fields', Q. J. Math. 5 (1934), 150–160.
- [6] J. Hoffstein and N. Jochnowitz, 'On Artin's conjecture and the class number of certain *CM* fields, I', *Duke Math. J.* **59** (1989), 553–563.
- [7] Y. Ihara, 'On the Euler–Kronecker constants of global fields and primes with small norms', in: *Algebraic Geometry and Number Theory*, Progress in Mathematics, 253 (ed. V. Ginzberg) (Birkhäuser, Boston, 2006), 407–451.
- [8] J. C. Lagarias and A. M. Odlyzko, 'Effective versions of the Chebotarev density theorem', in: *Algebraic Number Fields: L-Functions and Galois properties*, Proceedings of a Symposium, Durham University, Durham, 1975 (ed. A. Fröhlich) (Academic Press, London, 1977), 409–464.
- [9] S. R. Louboutin, 'Explicit upper bounds for residues of Dedekind zeta functions and values of *L*-functions at s = 1, and explicit lower bounds for relative class numbers of CM-fields', *Canad. J. Math.* 53 (2001), 1194–1222.
- [10] V. K. Murty, 'Class numbers of CM-fields with solvable normal closure', Compos. Math. 127 (2001), 273–287.
- [11] A. M. Odlyzko, 'Some analytic estimates of class numbers and discriminants', *Invent. Math.* 29 (1975), 275–286.
- [12] C. L. Siegel, 'Über die Classenzahl quadratischer Zahlkörper', Acta Arith. 1 (1935), 83-86.
- [13] H. M. Stark, 'Some effective cases of the Brauer-Siegel theorem', *Invent. Math.* 23 (1974), 135–152.
- [14] M. A. Tsfasman and S. G. Vlådut, 'Infinite global fields and the generalized Brauer-Siegel theorem', *Mosc. Math. J.* 2 (2002), 329–402.
- [15] A. Zykin, 'The Brauer–Siegel and Tsfasman–Vlädut theorems for almost normal extensions of number fields', Mosc. Math. J. 5 (2005), 961–968.

PENG-JIE WONG, National Center for Theoretical Sciences, No. 1, Sec. 4, Roosevelt Rd., Taipei City, Taiwan e-mail: pengjie.wong@ncts.tw